

Simulating expansions without expansions

A propos de la simulation des expansions sans expansions

Roberto Di Cosmo*

Delia Kesner §

Abstract

We add extensional equalities for the functional and product types to the typed λ -calculus with not only products and terminal object, but also sums and bounded recursion (a version of recursion that does not allow recursive calls of infinite length). We provide a confluent and strongly normalizing (thus decidable) rewriting system for the calculus, that stays confluent when allowing unbounded recursion. For that, we turn the extensional equalities into *expansion* rules, and not into contractions as is done traditionally. We first prove the calculus to be weakly confluent, which is a more complex and interesting task than for the usual λ -calculus. Then we provide an effective mechanism to simulate expansions without expansion rules, so that the strong normalization of the calculus can be derived from that of the underlying, traditional, non extensional system. These results give us the confluence of the full calculus, but we also show how to deduce confluence directly from our simulation technique, without the weak confluence property.

Résumé

On ajoute des égalités extensionnelles pour les types flèche et produit au λ -calcul typé avec produits, sommes, objet terminal et récursivité limitée (une forme de récursivité qui ne permet pas des appels récursifs de longueur infinie). On fournit un système de réécriture confluent et fortement normalisable (donc décidable), qui reste confluent quand l'on permet la récursivité illimitée. Pour cela, on transforme les égalités extensionnelles en règles d'*expansion*, interprétation différente de la traditionnelle qui les oriente comme des contractions. On prouve d'abord la confluence faible du calcul, ce qui est plus difficile et intéressant que pour le λ -calcul usuel. Ensuite, on donne une méthode effective pour simuler les expansions sans les règles d'expansions, de telle façon que la normalisation forte du calcul soit réduite à celle du système non extensionnel traditionnel sous-jacent. Ces résultats ainsi obtenus nous permettent de dériver la confluence du calcul entier, mais on montre aussi comment déduire directement la propriété de confluence à partir de la technique de simulation, sans avoir recours à la confluence faible.

*DMI-LIENS (CNRS URA 1347) Ecole Normale Supérieure - 45, Rue d'Ulm - 75230 Paris France
E-mail:dicosmo@dm.ens.fr

§INRIA Rocquencourt - Domaine de Voluceau, BP 105 - 78153 Le Chesnay Cedex, France and
CNRS and LRI - Bât 490, Université de Paris-Sud - 91405 Orsay Cedex, France
E-mail:kesner@margaux.inria.fr

Contents

1	Introduction	2
2	Survey	2
2.1	Our work	4
3	The Calculus	4
3.1	Types and Terms	4
3.2	Equality	5
3.3	The confluent rewriting system	6
3.4	Influential Positions	7
3.5	Adequacy of expansions for extensional equalities	8
3.6	Basic Properties of the Calculus	9
4	Weak Confluence	11
4.1	Some difficulties	11
4.2	Solving Critical Pairs	11
4.3	From Solved Critical Pairs to Full Weak Confluence	23
5	Strong Normalization	26
5.1	Simulating Expansions without Expansions	27
5.2	Strong Normalization of the Full Calculus	42
6	Strong Normalization via stability	42
6.1	Stability	43
6.2	Properties of stability	44
6.3	Products, sums and basic recursion	45
6.4	Abstraction and recursion	47
7	Strong Normalization via reducibility	49
7.1	Reducibility	50
7.2	Properties of reducibility	50
7.2.1	Atomic types	50
7.2.2	Product types	51
7.2.3	Arrow types	51
7.2.4	Sum types	51
7.3	Reducibility theorem	52
8	Confluence of the Full Calculus	55
9	Adding weak extensionality for the sum type	56
10	Conclusion and Future Work	57

1 Introduction

Over the past years there has been a growing interest in the properties of λ -calculus extended with various different type constructors, in particular pairs and sums, used to represent common data types. For these type constructors it is customary to provide a set of equalities that are then turned into computation rules: this is the case, for example, of the elimination rules for pairs:

$$(\pi_1) \quad \pi_1(\langle M, N \rangle) = M \quad (\pi_2) \quad \pi_2(\langle M, N \rangle) = N$$

They tell us how to operationally *compute* with objects of these types: if we have a pair $\langle M, N \rangle$, then we can decompose it to access its first or second component.

There is anyway something else that one likes to do with λ -calculus, besides using λ -terms as programs to be computed: one would like to *reason* about programs, to prove that they enjoy certain properties. Here is where extensional equalities come into play. In the case of functions, for example, since the only operational way to *use* a function is to apply it to an argument, we do not really want to consider a term M of function type different from the term $\lambda x.Mx$ where x does not occur free in M : both terms, when applied to an argument N , give the same result MN . Similarly for pairs, the only operational way to *use* a pair is by projecting out the first or the second component, so we do not want to consider a term M of product type different from the term $\langle \pi_1(M), \pi_2(M) \rangle$: the result of accessing any of these two terms via a first or second projection is the same term $\pi_1(M)$ or $\pi_2(M)$.

These facts can be incorporated in the calculus in the form of equalities, that one can read in at least two different ways:

- *an operational way*: these equalities just state possible *optimizations* of a program. Since a term $\langle \pi_1(M), \pi_2$

With the previous experience of the η rule, it is easy to understand how, at that time, most of the people thought that the right way to turn such an equality into a rewrite rule was also from left to right, as a contraction. But in 1980, J.W. Klop discovered [Klo80] that, if added to the usual confluent rewrite rules for pure λ -calculus, this interpretation of *SP* breaks confluence¹.

Anyway, this first negative result was shortly after mitigated in [Pot81] for the simply typed λ -calculus with η and *SP* contractions, by providing a first proof of confluence and strong normalization, later on simplified in different ways (see [Tro86] or [GLT90], for example). From then on, the contraction rule for *SP* was not considered harmful in a typed framework, until the seminal work by Lambek and Scott [LS86]. There, the decision problem of the equational theory of Cartesian Closed Categories (*ccc*'s) is solved using a particular typed λ -calculus equipped with not only η and *SP* equalities, but also with a special type \mathbf{T} representing the *terminal object* of the *ccc*'s². This distinguished atomic type comes with a further extensional axiom asserting that there is exactly one term $*$ of type \mathbf{T} :

$$(Top) \quad M : \mathbf{T} = *$$

Now, the type \mathbf{T} has the bad property of destroying confluence, if the extensional equalities η and *SP* are turned into contraction rules: the following are the critical pairs that arise immediately, as first pointed out by Obtulowicz, (see [LS86]):

$$\begin{array}{ccc} \langle \pi_1(x), \pi_2(x) \rangle \Rightarrow_{SP} x & & \langle \pi_1(x), \pi_2(x) \rangle \Rightarrow_{SP} x \\ \Downarrow^{Top} & & \Downarrow^{Top} \\ \langle *, \pi_2(x) \rangle & & \langle \pi_1(x), * \rangle \\ \\ (\lambda x : \mathbf{T}. Mx) : \mathbf{T} \rightarrow A \Rightarrow_{\eta} M & & (\lambda x : A. Mx) : A \rightarrow \mathbf{T} \Rightarrow_{\eta} M \\ \Downarrow_{Top} & & \Downarrow_{Top} \\ (\lambda x : \mathbf{T}. M*) : \mathbf{T} \rightarrow A & & (\lambda x : A. *) : A \rightarrow \mathbf{T} \end{array}$$

It is indeed possible, but not easy, to extend the contractive reduction system in order to recover confluence. A first step towards such a confluent system was taken by Poigné and Voss, who were not inspired by category theory, but by the implementation of algebraic data types [PV87]. In their paper, they study a calculus that includes $\lambda^1\beta\eta\pi*$, and notice that to solve the previous critical pairs one needs to add an infinite number of reduction rules (that can be anyway finitely described). Then confluence of such an extended system can be proved by showing weak confluence and strong normalization. Unfortunately, the critical pair for $(\lambda x : A. Mx) : \mathbf{T} \rightarrow A$ is missing there, and the strong normalization proof is incomplete.

More recently, Curien and the first author got interested in a polymorphic extension of $\lambda^1\beta\eta\pi*$, that arose in the study of the theory of object oriented programming and of isomorphisms of types [CDC91]. They give a complete (infinite) set of reduction rules for the calculus, which is proved confluent using just weak confluence, weak normalization and some additional properties.

Meanwhile, in the field of proof theory, Prawitz was suggesting [Pra71] to turn these extensional equalities into *expansion* rules, rather than contractions. Building on such ideas, but motivated by the study of coherence problems in category theory, Mints gives a first faulty proof that in the typed framework *expansion rules*, if handled with care, are weakly normalizing and preserve confluence of the typed calculus [Min79]³.

This idea of using expansion rules seems to have passed unnoticed for a long time, even if the so called η -long normal forms were well known and used in the study of higher order unification problems [Hue76]: only in these last years there has been a renewed interest in expansion rules. In recent work [Jay92], still motivated by category theoretic investigation, Jay explores a simply typed

¹See [Bar84], p. 403-409 for a short history and references.

²This is the *Unit* type in languages like ML.

³The same idea is present in [Min77].

λ -calculus with js \mathbf{T} and a natural number type \mathbf{N} as baeype, ewip with an inm-
 binator forms of type \mathbf{N} . He itrowcpansn rwl fr η and SP hat araclhe sme
 ls i prv an srnl

normlg.atcorher iloheotciatci ofwbr
 oriinal proof b Mis sowignflwec and wk nomal
 malizatcn, and [ka93] wher an itcereig diieand-
 same propere
 atci (cbwt nt srn nraltin.
 r prof of an srn nr
 cnqwer approach ipso pvehe

21 Owork

The presepaper isied b alhe priws wrs, bwt e
 wsepansin rwlso proie anflweteriig systcem
 prodwcs andermial objc, bwtlowms and recwrsn.
 of resie ssem where recwrsn i bowndedcrecrsi
 whih i proo be wklnflwetand stcrogl nomal
 orheype λ -lih ntn
 hi rele fmhen
 el of ifiniiegth ar nt alw,
 ng.

Wowhat song noraltci ofhe fl systcemn be re
 witchowtpansin rwl, fr whihhe tcradiial tcechni
 redwcin i thel(c)-0.607069uls wihpansns cn be *smlae*
 tchel(c)-0.607069uls wihowt epansns. Iwrs owthatchi re
 alshenflwence propery ashown i sectcn 8.
 dwcohat ofhe sse
 wes cn be ws (cwiw pofin
 at purpose, w swhat anyn se
 b a no-empy rinu i
 l i powfl enuh tco pvecl

Siche redwcin wih epansn rwl no a cngrwenc
 hold frhe w knonype λ -cllavo be rermwltce ihe epannr fine
 differeny as w wi s iecin 4. Frhi reon w bel
 rwle dereo be sdid mwcore crlo wi
 weaknflwence: thi wi lds tco wncanf tche esn
 , seea fnealpihat
 or i
 ie tchattheseihpnn
 ndeak tcheasf pvi dcl
 tcil ftcwresf thiin

W itrowc now tche clsnd is rewcinystcem i sec]TJ283.9230Td[(tc)-0.533549in 3,hen w ieiathe
 secin 4)n srn nraltin
 tc27.5517(w)28.19629o dientcys an fil ihe
 f tcehniwes

i paper. Therr twen
 nbownded rec]TJ283.6080Td[(w)-0.555882(r)-1.06599(s)-0.7743789in, tcha der ihe tce

⁴T stcnr tc tcmbljectnsror tc

Uni tc29.0453(y)-1.22222(p)-29.94899enanwgesik M.

$$\overline{\Gamma \vdash * : \mathbf{T}}$$

$\frac{}{x_1 : A_1, \dots, x_n : A_n \vdash x_i : A_i} \quad 1 \leq i \leq n \quad \text{and the } x_i\text{'s are pairwise distinct}$

$$\frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash \lambda x : A. M : A \rightarrow B}$$

$$\frac{\Gamma \vdash M : A \rightarrow B \quad \Gamma \vdash N : A}{\Gamma \vdash (MN) : B}$$

$$\frac{\Gamma \vdash M : A \quad \Gamma \vdash N : B}{\Gamma \vdash \langle M, N \rangle : A \times B}$$

$$\frac{\Gamma \vdash M : B_1 \times B_2}{\Gamma \vdash \pi_i(M) : B_i} \quad i = 1, 2$$

$$\frac{\Gamma \vdash M : B_i}{\Gamma \vdash in_{B_1+B_2}^i(M) : B_1 + B_2} \quad i = 1, 2$$

$$\frac{\Gamma \vdash P : A_1 + A_2 \quad \Gamma \vdash M_i : A_i \rightarrow D}{\Gamma \vdash Case(P, M_1, M_2) : D}$$

$$\frac{\Gamma, x : A \vdash M : A}{\Gamma \vdash (rec \ x : A. M)^i : A} \quad i \geq 0$$

We may omit types of variables in λ -abstractions when they are clear from the context writing $\lambda y. M$ instead of $\lambda y : C. M$.

Notation 3.1 (Free variables, substitutions) *The set of free variables of a term M will be noted $FV(M)$. It can be defined inductively as follows:*

$$\begin{aligned} FV(*) &= \emptyset \\ FV(x) &= \{x\} \\ FV(O_A) &= FV(M) \\ FV(MN) &= FV(M) + FV(N) \\ FV(\langle M, N \rangle) &= FV(M) + FV(N) \\ FV(\lambda x : A. M) &= FV(M) - \{x\} \\ FV((rec \ x : A. M)^i) &= FV(M) - \{x\} \\ FV(in_C^1(M)) &= FV(M) \\ FV(in_C^2(M)) &= FV(M) \\ FV(\pi_1(M)) &= FV(M) \\ FV(\pi_2(M)) &= FV(M) \\ FV(Case(P, M, N)) &= FV(P) + FV(M) + FV(N) \end{aligned}$$

We write $[N_1, \dots, N_n/x_1, \dots, x_n]$ (often abbreviated $[\overline{N}/\overline{x}]$) for the typed substitution mapping each variable $x_i : A_i$ to a term $N_i : A_i$. We write $M[\overline{N}/\overline{x}]$ for the term M where each variable x_i free in M is replaced by N_i .

3.2 Equality

Besides the usual identification of terms up to α conversion (i.e. renaming of bound variables), our calculus is equipped with the following equalities between terms.

$$\begin{array}{lll}
(\beta) & (\lambda x : A.M)N & = M[N/x] \\
(\pi_1) & \pi_1(\langle M_1, M_2 \rangle) & = M_1 \\
(\pi_2) & \pi_2(\langle M_1, M_2 \rangle) & = M_2 \\
(\rho) & \mathit{Case}(in_C^1(R), M_1, M_2) & = M_1 R \\
& \mathit{Case}(in_C^2(R), M_1, M_2) & = M_2 R \\
(\mathit{rec}) & (\mathit{rec } y : C.M)^{i+1} & = M[(\mathit{rec } y : C.M)^i/y] \\
(\eta) & \lambda x : A.Mx & = M \text{ if } \begin{cases} x \notin FV(M) \\ M : A \rightarrow B \end{cases} \\
(\delta) & \langle \pi_1(M), \pi_2(M) \rangle & = M \text{ if } M : A \times B \\
(\mathit{Top}) & M & = * \text{ if } M : \mathbf{T}
\end{array}$$

The index i that is attached to each rec term is a *bound* on the depth of the recursive calls that can originate from it. With such a bound, it is possible to insure the strong normalization of the associated reduction system.

The unbounded system is obtained from the bounded one by simply erasing all the bound indexes from the formation and equality rules (and the associated reduction rules). As we will show later, the bounded system can simulate any finite reduction of the unbounded system, and this fact will make it easy to extend the confluence result for the bounded system to the unbounded one. For simplicity, we will explicitly note the bound index only when necessary, dropping it whenever the properties we discuss hold in both systems.

3.3 The confluent rewriting system

The non extensional equality rules and the rule for \mathbf{T} can be turned into a confluent rewriting system by orienting them from left to right, as follows

$$\begin{array}{lll}
(\beta) & (\lambda x : A.M)N & \longrightarrow M[N/x] \\
(\pi_i) & \pi_i(\langle M_1, M_2 \rangle) & \longrightarrow M_i, \text{ for } i = 1, 2 \\
(\rho) & \mathit{Case}(in_C^i(R), M_1, M_2) & \longrightarrow M_i R, \text{ for } i = 1, 2 \\
(\mathit{rec}) & (\mathit{rec } y : C.M)^{i+1} & \longrightarrow M[(\mathit{rec } y : C.M)^i/y], \text{ for } i \geq 0 \\
(\mathit{Top}) & M & \longrightarrow * \text{ if } M : \mathbf{T} \text{ and } M \neq *
\end{array}$$

But when we want to turn the extensional equalities for functions and pairs into expansions, as explained very clearly by Jay, we must be careful to avoid the following reduction loops:

$$\begin{array}{lll}
\lambda x.M & \rightsquigarrow & \lambda y.(\lambda x.M)y & \rightsquigarrow & \lambda y.M[y/x] =_\alpha \lambda x.M \\
\langle M, N \rangle & \rightsquigarrow & \langle \pi_1(\langle M, N \rangle), \pi_2(\langle M, N \rangle) \rangle & \rightsquigarrow & \langle M, N \rangle \\
MN & \rightsquigarrow & (\lambda x.Mx)N & \rightsquigarrow & MN \\
\pi_i(P) & \rightsquigarrow & \pi_i(\langle \pi_1(P), \pi_2(P) \rangle) & \rightsquigarrow & \pi_i(P)
\end{array}$$

To break the first two loops we must disallow expansions of terms that are already λ -abstractions or pairs:

$$\begin{array}{ll}
(\eta) & M \longrightarrow \lambda x : A.Mx \text{ if } \begin{cases} x \notin FV(M) \\ M : A \rightarrow B \text{ and } M \text{ is not a } \lambda\text{-abstraction} \end{cases} \\
(\delta) & M \longrightarrow \langle \pi_1(M), \pi_2(M) \rangle \text{ if } \begin{cases} M : A \times B \text{ and } M \text{ is not a pair} \end{cases}
\end{array}$$

But this is not enough: to break the last two loops we must also forbid the η expansion of a term in a context where this term is applied to an argument, and δ expansion of a term when such a term is the argument of a projection. This means that we cannot define the one-step reduction relation \Longrightarrow on terms as the least congruence on terms containing the above reductions \longrightarrow , as is done usually. Instead, one defines formally $M \Longrightarrow M'$ starting from \longrightarrow by induction on the structure of the term. The definition is the same as a congruence closure but for the two last cases.

We will write $M \xrightarrow{\gamma_1 \dots \gamma_n} M'$ if $M \xrightarrow{\gamma_i} M'$, for some i and $\xrightarrow{\gamma}$ stands for a \longrightarrow step that is not a γ step. The one-step reduction relation between terms, denoted \Longrightarrow is defined as follows:

Definition 3.2 (One-step reduction)

- If $M \longrightarrow M'$, then $M \Longrightarrow M'$
- If $M \Longrightarrow M'$, then $(\text{rec } x : A.M)^i \Longrightarrow (\text{rec } x : A.M')^i$
 $\text{Case}(M, N, O) \Longrightarrow \text{Case}(M', N, O)$ $\text{in}_C^1(M) \Longrightarrow \text{in}_C^1(M')$ $\langle M, N \rangle \Longrightarrow \langle M', N \rangle$
 $\text{Case}(N, M, O) \Longrightarrow \text{Case}(N, M', O)$ $\text{in}_C^2(M) \Longrightarrow \text{in}_C^2(M')$ $\langle N, M \rangle \Longrightarrow \langle N, M' \rangle$
 $\text{Case}(N, O, M) \Longrightarrow \text{Case}(N, O, M')$ $\lambda x : A.M \Longrightarrow \lambda x : A.M'$ $NM \Longrightarrow NM'$
- If $M \Longrightarrow M'$ but $M \xrightarrow{\eta} M'$, then $MN \Longrightarrow M'N$
- If $M \Longrightarrow M'$ but $M \xrightarrow{\delta} M'$, then $\pi_i(M) \Longrightarrow \pi_i(M')$ for $i = 1, 2$

Notation 3.3 The transitive and the reflexive transitive closure of \Longrightarrow are noted \Longrightarrow^+ and \Longrightarrow^* respectively. Similarly we define \Longrightarrow^δ , $\Longrightarrow^{\delta+}$ and $\Longrightarrow^{\delta*}$ for the unbounded system.

We will use some standard notions from the theory of rewriting system, such as redex, normal form, confluence, weak confluence, strong normalization, etc, without explicitly redefining them here.

3.4 Influential Positions

It is also useful to define a notion of *influential positions* of a term: informally, a position in a term is *influential* if it prevents an expansion rule from being applied at the root of the subterm found at that position. For example, M occurs at an influential position in the term MN , as η expansion is forbidden on M , no matters if it is a λ -abstraction or not. Obviously, a position in a term can be influential for η or for δ , but not for both.

Formally, we define the set $\mathcal{P}(M)$ of *positions* of a term M and we distinguish two subsets of it: the set of influential positions for η , denoted $\mathcal{IP}_\eta(M)$, that prevent the η expansion rule at a subterm appearing at position $u \in \mathcal{IP}_\eta(M)$ and the set of influential positions for δ , denoted $\mathcal{IP}_\delta(M)$, that prevent the δ expansion rule at a subterm appearing at position $u \in \mathcal{IP}_\delta(M)$. The concatenation of the positions u and v is denoted $u.v$ and the concatenation of the position u with the set of positions \mathcal{S} is defined as $\{u.v \mid v \in \mathcal{S}\}$. When $\mathcal{S} = \emptyset$, $u.\mathcal{S} = \emptyset$. Formally:

$$\begin{aligned}
\mathcal{P}(\ast) &= \{\epsilon\} \\
\mathcal{P}(x) &= \{\epsilon\} \\
\mathcal{P}(M_1M_2) &= \{\epsilon\} + 1.\mathcal{P}(M_1) + 2.\mathcal{P}(M_2) \\
\mathcal{P}(\langle M_1, M_2 \rangle) &= \{\epsilon\} + 1.\mathcal{P}(M_1) + 2.\mathcal{P}(M_2) \\
\mathcal{P}(\lambda x : A.M) &= \{\epsilon\} + \{1\} + 2.\mathcal{P}(M) \\
\mathcal{P}(\pi_1(M)) &= \{\epsilon\} + 1.\mathcal{P}(M) \\
\mathcal{P}(\pi_2(M)) &= \{\epsilon\} + 1.\mathcal{P}(M) \\
\mathcal{P}(\text{Case}(P, M, N)) &= \{\epsilon\} + 1.\mathcal{P}(P) + 2.\mathcal{P}(M) + 3.\mathcal{P}(N) \\
\mathcal{P}((\text{rec } y : A.M)^i) &= \{\epsilon\} + \{1\} + 2.\mathcal{P}(M) \\
\mathcal{P}(\text{in}_C^1(M)) &= \{\epsilon\} + 1.\mathcal{P}(M) \\
\mathcal{P}(\text{in}_C^2(M)) &= \{\epsilon\} + 1.\mathcal{P}(M)
\end{aligned}$$

$$\begin{aligned}
\mathcal{IP}_\eta(\ast) &= \emptyset \\
\mathcal{IP}_\eta(x) &= \emptyset \\
\mathcal{IP}_\eta(M_1M_2) &= \{1\} + 1.\mathcal{IP}_\eta(M_1) + 2.\mathcal{IP}_\eta(M_2) \\
\mathcal{IP}_\eta(\langle M_1, M_2 \rangle) &= 1.\mathcal{IP}_\eta(M_1) + 2.\mathcal{IP}_\eta(M_2) \\
\mathcal{IP}_\eta(\lambda x : A.M) &= \{\epsilon\} + 2.\mathcal{IP}_\eta(M) \\
\mathcal{IP}_\eta(\pi_1(M)) &= 1.\mathcal{IP}_\eta(M) \\
\mathcal{IP}_\eta(\pi_2(M)) &= 1.\mathcal{IP}_\eta(M) \\
\mathcal{IP}_\eta(\text{Case}(P, M, N)) &= 1.\mathcal{IP}_\eta(P) + 2.\mathcal{IP}_\eta(M) + 3.\mathcal{IP}_\eta(N) \\
\mathcal{IP}_\eta((\text{rec } y : A.M)^i) &= 2.\mathcal{IP}_\eta(M) \\
\mathcal{IP}_\eta(\text{in}_C^1(M)) &= 1.\mathcal{IP}_\eta(M) \\
\mathcal{IP}_\eta(\text{in}_C^2(M)) &= 1.\mathcal{IP}_\eta(M)
\end{aligned}$$

$$\begin{aligned}
\mathcal{IP}_\delta(*) &= \emptyset \\
\mathcal{IP}_\delta(x) &= \emptyset \\
\mathcal{IP}_\delta(M_1 M_2) &= 1.\mathcal{IP}_\delta(M_1) + 2.\mathcal{IP}_\delta(M_2) \\
\mathcal{IP}_\delta(\langle M_1, M_2 \rangle) &= \{\epsilon\} + 1.\mathcal{IP}_\delta(M_1) + 2.\mathcal{IP}_\delta(M_2) \\
\mathcal{IP}_\delta(\lambda x : A.M) &= 2.\mathcal{IP}_\delta(M) \\
\mathcal{IP}_\delta(\pi_1(M)) &= \{1\} + 1.\mathcal{IP}_\delta(M) \\
\mathcal{IP}_\delta(\pi_2(M)) &= \{1\} + 1.\mathcal{IP}_\delta(M) \\
\mathcal{IP}_\delta(\text{Case}(P, M, N)) &= 1.\mathcal{IP}_\delta(P) + 2.\mathcal{IP}_\delta(M) + 3.\mathcal{IP}_\delta(N) \\
\mathcal{IP}_\delta(\text{rec } y : A.M^i) &= 2.\mathcal{IP}_\delta(M) \\
\mathcal{IP}_\delta(\text{in}_C^1(M)) &= 1.\mathcal{IP}_\delta(M) \\
\mathcal{IP}_\delta(\text{in}_C^2(M)) &= 1.\mathcal{IP}_\delta(M)
\end{aligned}$$

For example:

$$\begin{aligned}
\mathcal{P}(\langle xy, \langle \pi_1(x), zx \rangle \rangle) &= \{\epsilon, 1, 2, 1.1, 1.2, 2.1, 2.2, 2.1.1, 2.2.1, 2.2.2\} \\
\mathcal{IP}_\eta(\langle xy, \langle \pi_1(x), zx \rangle \rangle) &= \{1.1, 2.2.1\} \\
\mathcal{IP}_\delta(\langle xy, \langle \pi_1(x), zx \rangle \rangle) &= \{\epsilon, 2, 2.1.1\}
\end{aligned}$$

In general, we will say that u is an influential position of a term M if $u \in \mathcal{IP}_\eta(M)$ or $u \in \mathcal{IP}_\delta(M)$

3.5 Adequacy of expansions for extensional equalities

First of all, it is necessary to show that the limitations imposed on the reduction system do not make us loose any valid equality. We will show that the reduction system just introduced really generates the equalities we defined for the calculus. This comes from the fact that the limitations imposed on the reductions are introduced exactly to avoid reduction loops.

Theorem 3.4 (\implies **generates** E) *The equality E and the reflexive, symmetric and transitive closure R of \implies are the same relation.*

Proof.

The fact that R is included in E is evident, as all the reductions rules are derived from the equality axioms by orienting and restricting them.

What we are left to show is $E \subseteq R$. It is enough to show that whenever $M = N$ comes from a single equality axiom, we can either rewrite M to N or N to M (since R is reflexive, symmetric and transitive, the other cases will follow trivially).

The basic idea of the proof is to associate to each of these equality steps a reduction step in R . This is done in the obvious way, except in the cases that would violate one of the restrictions imposed on the expansion rules, which we will solve using exactly the reduction loop that this restriction is supposed to prevent.

Here are the problematic cases and how to deal with them. We use the usual context notation $C[M]$ to indicate a particular occurrence of a subterm M of interest in the term $C[M]$.

- $C[\lambda x.M] =_\eta C[\lambda y.(\lambda x.M)y]$. We cannot associate an η reduction to this equality, as we cannot expand something that is already an abstraction. But we can associate to it a β reduction from $C[\lambda y.(\lambda x.M)y]$ to $C[\lambda y.M[y/x]] = C[\lambda x.M]$.
- $C[\langle M, N \rangle] =_\delta \langle \pi_1(\langle M, N \rangle), \pi_2(\langle M, N \rangle) \rangle$. We cannot expand something that is already a pair, but we can use the π_i 's reduction from $\langle \pi_1(\langle M, N \rangle), \pi_2(\langle M, N \rangle) \rangle$ to $C[\langle M, N \rangle]$.
- $C[MN] =_\eta C[(\lambda x.Mx)N]$. Here we cannot expand M , which is in an influential position, but again we can use β to go from $C[(\lambda x.Mx)N]$ to $C[MN]$ (recall that $x \notin \text{FV}(M)$).
- $C[\pi_i(P)] =_\delta C[\pi_i(\langle \pi_1(P)$

3.6 Basic Properties of the Calculus

The following two lemmas are used to show that this calculus enjoys the subject reduction property, which guarantees that reductions preserve types.

Lemma 3.5 *If $\Gamma, x : A \vdash M : C$ and $x \notin FV(M)$, then $\Gamma \vdash M : C$.*

Proof. By induction on M . \square

Lemma 3.6 *If $\Gamma, x : A \vdash M : C$ and $\Gamma \vdash N : A$, then $\Gamma \vdash M[N/x] : C$.*

Proof. We show the property by induction on the structure of M

- $M \equiv *$. We have $*[N/x] = *$ and $\Gamma \vdash * : \mathbf{T}$ is an axiom.
- $M \equiv x$. We have $x[N/x] = N$ and the property trivially holds.
- $M \equiv y \neq x$. We have $y[N/x] = y$ and by lemma 3.5 the property holds.
- $M \equiv \lambda y : B.P$. We have $(\lambda y : B.P)[N/x] = \lambda y : B.(P[N/x])$. Since $\Gamma, x : A \vdash \lambda y : B.P : B \rightarrow D$ comes from $\Gamma, x : A, y : B \vdash P : D$ we have by induction hypothesis $\Gamma, y : B \vdash P[N/x] : D$ and thus $\Gamma \vdash \lambda y : B.(P[N/x]) : B \rightarrow D$.
- $M \equiv (M_1 M_2)$. We have $(M_1 M_2)[N/x] = (M_1[N/x])(M_2[N/x])$. Since $\Gamma, x : A \vdash (M_1 M_2) : C$ comes from $\Gamma, x : A \vdash M_1 : A \rightarrow C$ and $\Gamma, x : A \vdash M_2 : A$, by induction hypothesis $\Gamma \vdash M_1[N/x] : A \rightarrow C$ and $\Gamma \vdash M_2[N/x] : A$ and thus $\Gamma \vdash (M_1[N/x])(M_2[N/x]) : C$.
- $M \equiv \langle M_1, M_2 \rangle$. We have $\langle M_1, M_2 \rangle[N/x] = \langle M_1[N/x], M_2[N/x] \rangle$. Since $\Gamma, x : A \vdash \langle M_1, M_2 \rangle : C_1 \times C_2$ comes from $\Gamma, x : A \vdash M_1 : C_1$ and $\Gamma, x : A \vdash M_2 : C_2$, by induction hypothesis $\Gamma \vdash M_1[N/x] : C_1$ and $\Gamma \vdash M_2[N/x] : C_2$ and thus $\Gamma \vdash \langle M_1[N/x], M_2[N/x] \rangle : C_1 \times C_2$.
- $M \equiv in_{D_1+D_2}^i(P)$, for $i = 1, 2$. We have $in_{D_1+D_2}^i(P)[N/x] = in_{D_1+D_2}^i(P[N/x])$. Since $\Gamma, x : A \vdash in_{D_1+D_2}^i(P) : D_1 + D_2$ comes from $\Gamma, x : A \vdash P : D_i$, by induction hypothesis $\Gamma \vdash P[N/x] : D_i$ and thus $\Gamma \vdash in_{D_1+D_2}^i(P[N/x]) : D_1 + D_2$.
- $M \equiv \pi_i(P)$, for $i = 1, 2$. We have $\pi_i(P)[N/x] = \pi_i(P[N/x])$. Since $\Gamma, x : A \vdash \pi_i(P) : C_i$ comes from $\Gamma, x : A \vdash P : C_1 \times C_2$, by induction hypothesis $\Gamma \vdash P[N/x] : C_1 \times C_2$, and thus $\Gamma \vdash \pi_i(P[N/x]) : C_i$.
- $M \equiv Case(P, M_1, M_2)$. We have $Case(P, M_1, M_2)[N/x] = Case(P[N/x], M_1[N/x], M_2[N/x])$. $\Gamma, x : A \vdash Case(P, M_1, M_2) : C$ comes from $\Gamma, x : A \vdash P : B_1 + B_2$, $\Gamma, x : A \vdash M_1 : B_1 \rightarrow C$ and $\Gamma, x : A \vdash M_2 : B_2 \rightarrow C$. By induction hypothesis $\Gamma \vdash P[N/x] : B_1 + B_2$, $\Gamma \vdash M_1[N/x] : B_1 \rightarrow C$ and $\Gamma \vdash M_2[N/x] : B_2 \rightarrow C$ and thus $\Gamma \vdash Case(P[N/x], M_1[N/x], M_2[N/x]) : C$.
- $M \equiv (rec\ y : C.P)^i$. We have $(rec\ y : C.P)^i[N/x] = (rec\ y : C.P[N/x])^i$. Since $\Gamma, x : A \vdash (rec\ y : C.P)^i : C$ comes from $\Gamma, x : A, y : C \vdash P : C$, by induction hypothesis $\Gamma, y : C \vdash P[N/x] : C$ and thus $\Gamma \vdash (rec\ y : C.P[N/x])^i : C$.

\square

Proposition 3.7 (Subject Reduction) *If $\Gamma \vdash R : C$ and $R \Longrightarrow R'$, then $\Gamma \vdash R' : C$*

Proof. We proceed by cases. Let's see first the case of one external reduction step:

- $(\lambda x : A.M)N \xrightarrow{\beta} M[N/x]$. As $\Gamma \vdash (\lambda x : A.M)N : C$ comes from $\Gamma, x : A \vdash M : C$ and $\Gamma \vdash N : A$, then $\Gamma \vdash M[N/x] : C$ holds by lemma 3.6.
- $M \xrightarrow{\eta} \lambda x : A.Mx$. As $\Gamma \vdash M : A \rightarrow D$ and $\Gamma, x : A \vdash x : A$, then $\Gamma, x : A \vdash Mx : D$ and $\Gamma \vdash \lambda x : A.Mx : A \rightarrow D$.
- $M \xrightarrow{Top} *$ and $\Gamma \vdash * : \mathbf{T}$ is an axiom.
- $\pi_i(\langle M_1, M_2 \rangle) \xrightarrow{\pi_i} M_i$. We have $\Gamma \vdash \pi_i(\langle M_1, M_2 \rangle) : C_i$ if $\Gamma \vdash \langle M_1, M_2 \rangle : C_1 \times C_2$ and this holds if $\Gamma \vdash M_i : C_i$.
- $M \xrightarrow{\delta} \langle \pi_1(M), \pi_2(M) \rangle$. Since $\Gamma \vdash M : C_1 \times C_2$, then $\Gamma \vdash \pi_1(M) : C_1$ and $\Gamma \vdash \pi_2(M) : C_2$ and thus $\Gamma \vdash \langle \pi_1(M), \pi_2(M) \rangle : C_1 \times C_2$.

- $Case(in_{B_1+B_2}^i(P, M_1, M_2) \xrightarrow{\rho} M_i P)$. Since $\Gamma \vdash Case(in_{B_1+B_2}^i(P), M_1, M_2) : C$, then $\Gamma \vdash in_{B_1+B_2}^i(P) : B_1 + B_2$, $\Gamma \vdash M_1 : B_1 \rightarrow C$ and $\Gamma \vdash M_2 : B_2 \rightarrow C$. Then we have $\Gamma \vdash P : B_i$ and $\Gamma \vdash M_i : B_i \rightarrow C$ and therefore $\Gamma \vdash M_i P : C$.
- $(rec\ y : C.M)^{i+1} \xrightarrow{rec} M[(rec\ y : C.M)^i/y]$. Since $\Gamma \vdash (rec\ y : C.M)^{i+1} : C$, then $\Gamma, y : C \vdash M : C$ and $\Gamma \vdash (rec\ y : C.M)^i : C$. Applying lemma 3.6 $\Gamma \vdash M[(rec\ y : C.M)^i/y] : C$.

Now, let's see the case of one internal reduction step.

- If $MN \Longrightarrow M'N$, where $M \Longrightarrow M'$. We have $\Gamma \vdash MN : C$ if $\Gamma \vdash M : A \rightarrow C$ and $\Gamma \vdash N : A$. By induction hypothesis $\Gamma \vdash M' : A \rightarrow C$ and thus $\Gamma \vdash M'N : C$.
- If $MN \Longrightarrow MN'$, where $N \Longrightarrow N'$. We have $\Gamma \vdash MN : C$ if $\Gamma \vdash M : A \rightarrow C$ and $\Gamma \vdash N : A$. By induction hypothesis $\Gamma \vdash N' : A$ and thus $\Gamma \vdash MN' : C$.
- $\lambda x : A.M \Longrightarrow \lambda x : A.M'$, where $M \Longrightarrow M'$. We have $\Gamma \vdash \lambda x : A.M : A \rightarrow B$ if $\Gamma, x : A \vdash M : B$. By induction hypothesis $\Gamma, x : A \vdash M' : B$ and thus $\Gamma \vdash \lambda x : A.M' : A \rightarrow B$.
- $\langle M, N \rangle \Longrightarrow \langle M', N \rangle$ if $M \Longrightarrow M'$. We have $\Gamma \vdash \langle M, N \rangle : C_1 \times C_2$ if $\Gamma \vdash M : C_1$ and $\Gamma \vdash N : C_2$. By induction hypothesis $\Gamma \vdash M' : C_1$ and thus $\Gamma \vdash \langle M', N \rangle : C_1 \times C_2$.
- $\langle M, N \rangle \Longrightarrow \langle M, N' \rangle$ if $N \Longrightarrow N'$. We have $\Gamma \vdash \langle M, N \rangle : C_1 \times C_2$ if $\Gamma \vdash M : C_1$ and $\Gamma \vdash N : C_2$. By induction hypothesis $\Gamma \vdash N' : C_2$ and thus $\Gamma \vdash \langle M, N' \rangle : C_1 \times C_2$.
- $\pi_i(M) \Longrightarrow \pi_i(M')$, for $i = 1, 2$ where $M \Longrightarrow M'$. We have $\Gamma \vdash \pi_i(M) : C_i$ if $\Gamma \vdash M : C_1 \times C_2$. By induction hypothesis $\Gamma \vdash M' : C_1 \times C_2$ and thus $\Gamma \vdash \pi_i(M') : C_i$.
- $in_{D_1+D_2}^i(M) \Longrightarrow in_{D_1+D_2}^i(M')$, for $i = 1, 2$ where $M \Longrightarrow M'$. We have $\Gamma \vdash in_{D_1+D_2}^i(M) : D_1 + D_2$ if $\Gamma \vdash M : D_i$. By induction hypothesis $\Gamma \vdash M' : D_i$ and thus $\Gamma \vdash in_{D_1+D_2}^i(M') : D_1 + D_2$.
- $Case(P, M, N) \Longrightarrow Case(P', M, N)$, where $P \Longrightarrow P'$. We have $\Gamma \vdash Case(P, M, N) : C$ if $\Gamma \vdash P : A + B$, $\Gamma \vdash M : A \rightarrow C$ and $\Gamma \vdash N : B \rightarrow C$. By induction hypothesis $\Gamma \vdash P' : A + B$ and thus $\Gamma \vdash Case(P', M, N) : C$.
- $Case(P, M, N) \Longrightarrow Case(P, M', N)$, where $M \Longrightarrow M'$. We have $\Gamma \vdash Case(P, M, N) : C$ if $\Gamma \vdash P : A + B$, $\Gamma \vdash M : A \rightarrow C$ and $\Gamma \vdash N : B \rightarrow C$. By induction hypothesis $\Gamma \vdash M' : A \rightarrow C$ and thus $\Gamma \vdash Case(P, M', N) : C$.
- $Case(P, M, N) \Longrightarrow Case(P, M, N')$, where $N \Longrightarrow N'$. We have $\Gamma \vdash Case(P, M, N) : C$ if $\Gamma \vdash P : A + B$, $\Gamma \vdash M : A \rightarrow C$ and $\Gamma \vdash N : B \rightarrow C$. By induction hypothesis $\Gamma \vdash N' : B \rightarrow C$ and thus $\Gamma \vdash Case(P, M, N') : C$.
- $(rec\ x : C.M)^i \Longrightarrow (rec\ x : C.M')^i$, where $M \Longrightarrow M'$. We have $\Gamma \vdash (rec\ x : C.M)^i : C$ if $\Gamma, x : C \vdash M : C$. By induction hypothesis $\Gamma, x : C \vdash M' : C$ and thus $\Gamma \vdash (rec\ x : C.M')^i : C$.

□

Another remarkable property of this calculus can be stated as follows:

Lemma 3.8 *If M is in normal form w.r.t. the system without the η , δ and Top rules and $M \xrightarrow{\eta, \delta, Top} R$, then R is in normal form w.r.t. the system without η , δ and Top .*

Proof. Suppose M has no β , π_i , ρ or rec redexes. Notice first that a ρ redex cannot be created in R as there is no way to introduce an in^i expression using the η , δ and Top rules. The same for rec . We are left with the following two cases:

- Suppose R has a β redex. Then $R \equiv C[(\lambda x.P)N]$ and since M contains no β redexes, we have necessarily $M \equiv C[SN]$, $P \equiv Sx$ and $S \xrightarrow{\eta} \lambda x.Sx$. But this is not possible because η expansions are not allowed on terms appearing in influential positions for η .
- Suppose R has a π_i redex. Then $R \equiv C[\pi_i(\langle M, N \rangle)]$ and since M contains no π_i 's redexes, we have necessarily $M \equiv C[\pi_i(T)]$, $M \equiv \pi_1(T)$, $N \equiv \pi_2(T)$ and $T \xrightarrow{\delta} \langle \pi_1(T), \pi_2(T) \rangle$. But this is not possible because δ expansions are not allowed on terms appearing in influential positions for δ .

□

Corollary 3.9 *If M is in normal form with respect to the system without the η , δ and Top rules, then the $\eta - \delta - \text{Top}$ normal form of M is in normal form.*

4 Weak Confluence

In this section we set off to prove that the reduction system proposed above is actually weakly confluent, i.e. that whenever $M' \leftarrow M \Rightarrow M''$ we can find a term M''' s.t. $M' \Rightarrow^* M''' \leftarrow^* M''$. The proof is fairly more complex here than in the case of λ -calculus where extensional equalities are interpreted as contractions, and this is due to the fact that the reduction relation \Rightarrow introduced above *is not a congruence on terms*.

4.1 Some difficulties

In particular, in the simply typed λ -calculus whenever $M \Rightarrow^* M'$ then $\pi_i(M) \Rightarrow^* \pi_i(M')$, and if also $N \Rightarrow^* N'$, then $MN \Rightarrow^* M'N'$, but this is no longer true now: indeed, we have $x : A \rightarrow B \Rightarrow \lambda z : A.xz$, but xN cannot reduce to $(\lambda z : A.xz)N$.

These properties still hold for those reduction sequences $M \Rightarrow^* M'$ that do not involve expansions at the root:

Remark 4.1

- Let $M \equiv M_0 \Rightarrow M_1 \Rightarrow \dots \Rightarrow M_{n-1} \Rightarrow M_n \equiv M'$ be a reduction sequence and let $N \Rightarrow^* N'$, where in the first reduction sequence there are no expansions applied at the root positions of the M_i 's. Then, $MN \Rightarrow^* M'N'$.
- Let $M \equiv M_0 \Rightarrow M_1 \Rightarrow \dots \Rightarrow M_{n-1} \Rightarrow M_n \equiv M'$ be a reduction sequence where none of the M_i 's is expanded at the root. Then $\pi_i(M) \Rightarrow^* \pi_i(M')$, for $i = 1, 2$.

Proof.

- Take the reduction $M \equiv M_0 \Rightarrow M_1 \Rightarrow \dots \Rightarrow M_{n-1} \Rightarrow M_n \equiv M'$: since there are no η expansions at the root we can form the reduction sequence $MN \Rightarrow M_1N \Rightarrow \dots \Rightarrow M_{n-1}N \Rightarrow M_nN \equiv M'N$. Then, we take the sequence $N \Rightarrow N_1 \Rightarrow \dots \Rightarrow N_{p-1} \Rightarrow N_p \equiv N'$ and we propagate it on the right of M' , i.e. $M'N \Rightarrow M'N_1 \Rightarrow \dots \Rightarrow M'N_{p-1} \Rightarrow M'N_p \equiv M'N'$.
- Take the reduction $M \Rightarrow M_1 \Rightarrow \dots \Rightarrow M_{n-1} \Rightarrow M_n \equiv M'$. Since there are no δ expansions at the root, we can form the reduction sequence $\pi_i(M) \Rightarrow \pi_i(M_1) \Rightarrow \dots \Rightarrow \pi_i(M_{n-1}) \Rightarrow \pi_i(M_n) \equiv \pi_i(M')$.

□

4.2 Solving Critical Pairs

In this calculus, it is no longer true that reduction is stable by substitution, as in the traditional λ -calculus:

Remark 4.2

If $P \Rightarrow P'$, $N \Rightarrow N'$, it is not true in general that $P[N/x] \Rightarrow^ P'[N/x]$ and $P[N/x] \Rightarrow^* P[N'/x]$.*

Indeed, $x : A \rightarrow B \Rightarrow \lambda z : A.xz$, but $x[\lambda y : A.w/x] = \lambda y : A.w$ cannot reduce in our system to $\lambda z : A.(\lambda y : A.w)z = \lambda z : A.xz[\lambda y : A.w/x]$, and $(yM)[x/y] = xM$ cannot reduce to $(\lambda z : A.xz)M = (yM)[\lambda z : A.xz/y]$.

We can prove some weaker properties: if $P \Rightarrow P'$, then $P[N/x]$ and $P'[N/x]$ have a common reduct (Lemma 4.5), and similarly $P[N/x]$ and $P[N'/x]$ when $N \Rightarrow N'$ (Lemma 4.6). This suffices for our purpose of proving weak confluence of the reduction system.

First of all it is useful to recall here a basic property of substitutions that do hold in our calculus.

Lemma 4.3 *If $x \neq y$ and $x \notin FV(L)$, then*

$$M[N/x][L/y] = M[L/y][N[L/y]/x]$$

Lemma 4.4 *If $P \xrightarrow{\eta, \delta, T \circ p} P'$, then $P[N/x] \Longrightarrow^* P'[N/x]$ or $P'[N/x] \Longrightarrow^* P[N/x]$. Moreover, if the expansion does not take place at the root of P , then there are no expansions at root positions in the sequences $P[N/x] \Longrightarrow^* P'[N/x]$ and $P'[N/x] \Longrightarrow^* P[N/x]$.*

Proof.

- $P \xrightarrow{\eta} \lambda z.Pz$. Then P is not a λ -abstraction, $P'[N/x] = \lambda z.P[N/x]z$ and there are two possible cases:

- If $P[N/x]$ is not a λ -abstraction, $P[N/x] \xrightarrow{\eta} \lambda z.P[N/x]z$ since P is of type \rightarrow and so $P[N/x]$ is also of type \rightarrow by lemma 3.6.
- If $P[N/x]$ is a λ -abstraction, then $P \equiv x$, $N \equiv \lambda y.N'$ and:
 $(\lambda z.xz)[\lambda y.N'/x] = \lambda z.(\lambda y.N')z \xrightarrow{\beta} \lambda z.(N'[z/y]) =_{\alpha} \lambda y.N' = x[\lambda y.N'/x]$.

- $P \xrightarrow{\delta} \langle \pi_1(P), \pi_2(P) \rangle$. Then P is not a pair, $P'[N/x] = \langle \pi_1(P[N/x]), \pi_2(P[N/x]) \rangle$ and there are two possible cases:

- If $P[N/x]$ is not a Pair, $P[N/x] \xrightarrow{\delta} \langle \pi_1(P[N/x]), \pi_2(P[N/x]) \rangle$ since P is of type \times and so $P[N/x]$ is also of type \times by lemma 3.6.
- If $P[N/x]$ is a pair, then $P \equiv x$ and $N \equiv \langle N_1, N_2 \rangle$ and:

$$\begin{aligned} & \langle \pi_1(x), \pi_2(x) \rangle [\langle N_1, N_2 \rangle / x] \\ &= \langle \pi_1(\langle N_1, N_2 \rangle), \pi_2(\langle N_1, N_2 \rangle) \rangle \end{aligned}$$

$$\begin{aligned} & \langle \pi_1(\langle N_1, N_2 \rangle), \pi_2(\langle N_1, N_2 \rangle) \rangle \end{aligned}$$

4. $P \equiv P_1 P_2$. There are several cases to consider:

- If $P \xrightarrow{\eta, \delta, Top} P'$ the property holds by lemma 4.4.
- If $P \equiv (\lambda z. P_3) P_2$, then $(\lambda z. P_3) P_2 \xrightarrow{\beta} P_3[N/x]$ and

$$(\lambda z. P_3) P_2[N/x] = (\lambda z. P_3[N/x]) P_2[N/x] \xrightarrow{\beta} P_3[N/x][P_2[N/x]/z] \stackrel{lemma 4.3}{=} P_3[P_2/z][N/x]$$

- If $P' \equiv P'_1 P_2$, where $P_1 \Longrightarrow P'_1$. By i.h. $P_1[N/x] \Longrightarrow^* P'_1[N/x]$ or $P'_1[N/x] \Longrightarrow^* P_1[N/x]$. Suppose the first case holds. Since $P_1 P_2 \Longrightarrow P'_1 P_2$, then $P_1 \xrightarrow{\eta} P'_1$ is not possible and by lemma 4.4 there are no η expansions at the root positions of the terms appearing in $P_1[N/x] \Longrightarrow^* P'_1[N/x]$. We have

$$(P_1 P_2)[N/x] = P_1[N/x] P_2[N/x] \Longrightarrow^* P'_1[N/x] P_2[N/x] = (P'_1 P_2)[N/x]$$

In the same way the property holds for the case $P'_1[N/x] \Longrightarrow^* P_1[N/x]$.

- If $P' \equiv P_1 P'_2$, where $P_2 \Longrightarrow P'_2$. By i.h. $P_2[N/x] \Longrightarrow^* P'_2[N/x]$ or $P'_2[N/x] \Longrightarrow^* P_2[N/x]$. Suppose the first case holds, then

$$(P_1 P_2)[N/x] = P_1[N/x] P_2[N/x] \Longrightarrow^* P_1[N/x] P'_2[N/x] = (P_1 P'_2)[N/x]$$

In the same way the property holds for the case $P'_2[N/x] \Longrightarrow^* P_2[N/x]$.

5. $P \equiv in^i_{C_1+C_2}(Q)$. Since $in^i_{C_1+C_2}(Q)$ is of type $C_1 + C_2$, then $P \xrightarrow{\eta, \delta, Top} P'$ is not possible. Therefore $P' \equiv in^i_{C_1+C_2}(Q')$, where $Q \Longrightarrow Q'$ and by induction hypothesis $Q[N/x] \Longrightarrow^* Q'[N/x]$ or $Q'[N/x] \Longrightarrow^* Q[N/x]$.

If the first case,

$$in^i_{C_1+C_2}(Q)[N/x] = in^i_{C_1+C_2}(Q[N/x]) \Longrightarrow^* in^i_{C_1+C_2}(Q'[N/x]) = in^i_{C_1+C_2}(Q')[N/x]$$

In the second

$$in^i_{C_1+C_2}(Q')[N/x] = in^i_{C_1+C_2}(Q'[N/x]) \Longrightarrow^* in^i_{C_1+C_2}(Q[N/x]) = in^i_{C_1+C_2}(Q)[N/x]$$

6. $P \equiv \langle P_1, P_2 \rangle$. Since $P \xrightarrow{\delta} P'$ is not possible because P is a pair, $P \xrightarrow{\eta} P'$ is neither possible because P is not of type \rightarrow , and $P \xrightarrow{Top} P'$ is neither possible because P is not of type \mathbf{T} , we have $P' \equiv \langle P'_1, P_2 \rangle$, where $P_1 \Longrightarrow P'_1$ or $P' \equiv \langle P_1, P'_2 \rangle$, where $P_2 \Longrightarrow P'_2$.

In the first case $P_1[N/x] \Longrightarrow^* P'_1[N/x]$ or $P'_1[N/x] \Longrightarrow^* P_1[N/x]$ hold by induction hypothesis and then either

$$\langle P_1, P_2 \rangle[N/x] = \langle P_1[N/x], P_2[N/x] \rangle \Longrightarrow^* \langle P'_1[N/x], P_2[N/x] \rangle = \langle P'_1, P_2 \rangle[N/x]$$

or

$$\langle P'_1, P_2 \rangle[N/x] \Longrightarrow^* \langle P_1, P_2 \rangle[N/x]$$

In the second case $P_2[N/x] \Longrightarrow^* P'_2[N/x]$ or $P'_2[N/x] \Longrightarrow^* P_2[N/x]$ hold by induction hypothesis and then

$$\langle P_1, P_2 \rangle[N/x] = \langle P_1[N/x], P_2[N/x] \rangle \Longrightarrow^* \langle P_1[N/x], P'_2[N/x] \rangle = \langle P_1, P'_2 \rangle[N/x]$$

or

$$\langle P_1, P'_2 \rangle[N/x] \Longrightarrow^* \langle P_1, P_2 \rangle[N/x]$$

7. $P \equiv \pi_i(Q)$. If $P \xrightarrow{\eta, \delta, Top} P'$ the property holds by lemma 4.4. If not, $P' \equiv \pi_i(Q')$, where $Q \Longrightarrow Q'$ and by induction hypothesis $Q[N/x] \Longrightarrow^* Q'[N/x]$ or $Q'[N/x] \Longrightarrow^* Q[N/x]$. Since $\pi_i(Q) \Longrightarrow \pi_i(Q')$, then $Q \rightarrow Q'$ is not a δ -expansion and thus there are no δ -expansions appearing at the root positions of the terms in $Q[N/x] \Longrightarrow^* Q'[N/x]$ and $Q'[N/x] \Longrightarrow^* Q[N/x]$ (first and second case resp.). Therefore either

$$\pi_i(Q)[N/x] = \pi_i(Q[N/x]) \Longrightarrow^* \pi_i(Q'[N/x]) = \pi_i(Q')[N/x]$$

or

$$\pi_i(Q')[N/x] \Longrightarrow^* \pi_i(Q)[N/x]$$

8. $P \equiv Case(Q, M_1, M_2)$. If $P \xrightarrow{\eta, \delta, Top} P'$ the property holds by lemma 4.4. If not, there are five possibilities:

- $Q \equiv in_{B_1+B_2}^i(R)$ and $Case(in_{B_1+B_2}^i(R), M_1, M_2) \xrightarrow{p} M_i R$. Since

$$Case(in_{B_1+B_2}^i(R), M_1, M_2)[N/x] = Case(in_{B_1+B_2}^i(R[N/x]), M_1[N/x], M_2[N/x])$$

and this last term reduces by a ρ -rule to $M_i[N/x]R[N/x] = (M_i R)[N/x]$ the property holds.

- $P' \equiv Case(Q', M_1, M_2)$, where $Q \Longrightarrow Q'$. By induction hypothesis $Q[N/x] \Longrightarrow^* Q'[N/x]$ or $Q'[N/x] \Longrightarrow^* Q[N/x]$.

In the first case

$$\begin{aligned} Case(Q, M_1, M_2)[N/x] &= Case(Q', M_1, M_2)[N/x] \\ &= Case(Q[N/x], M_1[N/x], M_2[N/x]) \Longrightarrow^* Case(Q'[N/x], M_1[N/x], M_2[N/x]) \end{aligned}$$

In the second case $Case(Q', M_1, M_2)[N/x] \Longrightarrow^* Case(Q, M_1, M_2)[N/x]$.

- $P' \equiv Case(Q, M'_1, M_2)$, where $M_1 \Longrightarrow M'_1$. By induction hypothesis $M_1[N/x] \Longrightarrow^* M'_1[N/x]$ or $M'_1[N/x] \Longrightarrow^* M_1[N/x]$.

In the first case

$$\begin{aligned} Case(Q, M_1, M_2)[N/x] &= Case(Q, M'_1, M_2)[N/x] \\ &= Case(Q[N/x], M_1[N/x], M_2[N/x]) \Longrightarrow^* Case(Q[N/x], M'_1[N/x], M_2[N/x]) \end{aligned}$$

In the second case $Case(Q, M'_1, M_2)[N/x] \Longrightarrow^* Case(Q, M_1, M_2)[N/x]$.

- $P' \equiv Case(Q, M_1, M'_2)$, where $M_2 \Longrightarrow M'_2$. By induction hypothesis $M_2[N/x] \Longrightarrow^* M'_2[N/x]$ or $M'_2[N/x] \Longrightarrow^* M_2[N/x]$.

In the first case

$$\begin{aligned} Case(Q, M_1, M_2)[N/x] &= Case(Q, M_1, M'_2)[N/x] \\ &= Case(Q[N/x], M_1[N/x], M_2[N/x]) \Longrightarrow^* Case(Q[N/x], M_1[N/x], M'_2[N/x]) \end{aligned}$$

In the second case $Case(Q, M_1, M'_2)[N/x] \Longrightarrow^* Case(Q, M_1, M_2)[N/x]$.

9. $P \equiv (rec\ y : B.Q)^i$. If $P \xrightarrow{\eta, \delta, Top} P'$ the property holds by lemma 4.4. If not there are two cases to consider:

- $P' \equiv Q[(rec\ y : B.Q)^{i-1}/y]$. We have $(rec\ y : B.Q)^i[N/x] = (rec\ y : B.Q[N/x])^i$ and this last term reduces to $Q[N/x][(rec\ y : B.Q[N/x])^{i-1}/y]$ that is equal to $Q[(rec\ y : B.Q)^{i-1}[N/x]/y][N/x] = Q[(rec\ y : B.Q)^{i-1}/y][N/x]$ by lemma 4.3.

- $P' \equiv (\text{rec } y : B.Q')^i$, where $Q \Longrightarrow Q'$. By induction hypothesis $Q[N/x] \Longrightarrow^* Q'[N/x]$ or $Q'[N/x] \Longrightarrow^* Q[N/x]$ and thus

$$(\text{rec } y : B.Q)^i[N/x] = (\text{rec } y : B.Q[N/x])^i \Longrightarrow^* (\text{rec } y : B.Q'[N/x])^i = (\text{rec } y : B.Q')^i[N/x]$$

or

$$(\text{rec } y : B.Q')^i[N/x] \Longrightarrow^* (\text{rec } y : B.Q)^i[N/x]$$

□

Lemma 4.6 (Substitution Lemma (ii))

If $N \xrightarrow{R} N'$, then $M[N/x] \Longrightarrow^* M'' * \leftarrow M[N'/x]$ for some term M'' . These reduction sequences contain expansions at the root only if $M \equiv x$ and R is an expansion applied at the root of N .

Proof. We will show that $M[N/x] \xrightarrow{R} M'' * \leftarrow M[N'/x]$ for some term M'' and that these reduction sequences contain expansions at the root only if $M \equiv x$ and R is an expansion applied at the root of N .

This is a very common lemma in the theory of λ -calculus, where the term M'' is always $M[N'/x]$ and the proof is straightforward context closure of the reduction R . Here the conditions imposed on the expansion rules make it necessary to state the lemma this way. Effectively, the only interesting cases of the proof are the ones for application and projections, where we cannot always apply context closure for the reduction R , and have to make some steps backwards from $M[N'/x]$ to $M[N/x]$.

Notice that every time the required reductions are built by context closure, there is no rule applied at the root and we state this fact here once for all. We proceed by induction on M :

- $M \equiv x$
 $M[N/x] = N \xrightarrow{R} N' = M[N'/x]$ (in this case our M'' is N')
- $M \equiv y \neq x$ or $M \equiv * : \mathbf{T}$
Then $M[N/x] = M = M[N'/x]$ (in this case our M'' is M)
- $M \equiv (M_1 M_2)$
We find by induction hypothesis terms M_1'' and M_2'' such that

$$M_1[N/x] \xrightarrow{R} M_1'' * \leftarrow M_1[N'/x], \text{ and } M_2[N/x] \xrightarrow{R} M_2'' * \leftarrow M_2[N'/x].$$

Here M_1 is in an influential position for η , so we have to be careful about the reductions occurring in $M_1[N/x] \xrightarrow{R} M_1'' * \leftarrow M_1[N'/x]$. We have the following cases:

- If $M_1 \neq x$, or R is not an expansion at the root of N , we know by inductive hypothesis that the reductions $M_1[N/x] \xrightarrow{R} M_1'' * \leftarrow M_1[N'/x]$ do not contain any expansions, and in particular no η rule, at the root position, so we can apply context closure for application and get

$$\begin{array}{ccc} (M_1 M_2)[N/x] & & (M_1 M_2)[N'/x] \\ = & & = \\ (M_1[N/x] M_2[N/x]) \xrightarrow{R} M_1'' * \leftarrow M_1[N'/x] & & (M_1[N'/x] M_2[N'/x]) \end{array}$$

- If $M_1 \equiv x$, and the expansion rule R is η at the root of N , then $N' \equiv \lambda z.Nz$ and we can close our diagram as follows

$$\begin{array}{ccc} (x[N/x] M_2[N/x]) & & (x[\lambda z.Nz/x] M_2[\lambda z.Nz/x]) \\ = & & = \\ (N M_2[N/x]) & & ((\lambda z.Nz) M_2[\lambda z.Nz/x]) \\ \Downarrow & \xleftarrow{\beta} & \Downarrow \\ (N M_2'') & & ((\lambda z.Nz) M_2'') \end{array}$$

Here, the vertical reductions are built by context closure, while the horizontal one is a β , so no expansion rule is applied at the root in the overall reduction sequence.

- $M \equiv \lambda z : A.M_1$

If $x \neq z$, the result follows from $(\lambda z : A.M_1)[N/x] = \lambda z : A.M_1 = (\lambda z : A.M_1)[N'/x]$. Otherwise, by induction hypothesis there is a term M_1'' such that $M_1[N/x] \xrightarrow{R}^* M_1'' * \leftarrow M_1[N'/x]$, so we can apply the context closure rule for abstraction and get that

$$\begin{array}{ccc} (\lambda z : A.M_1)[N/x] & & (\lambda z.M_1)[N'/x] \\ = & & = \\ (\lambda z : A.M_1[N/x]) \xrightarrow{R}^* \lambda z : A.M_1'' * \leftarrow & & (\lambda z : A.M_1[N'/x]) \end{array}$$

- $M \equiv \pi_i(M_1)$

We find by induction hypothesis a term M_1'' such that

$$M_1[N/x] \xrightarrow{R}^* M_1'' * \leftarrow M_1[N'/x].$$

Here M_1 is in an influential position for δ , so we have to be careful about the reductions occurring in $M_1[N/x] \xrightarrow{R}^* M_1'' * \leftarrow M_1[N'/x]$. We have the following cases:

- If $M_1 \not\equiv x$, or R is not an expansion at the root of N , we know by inductive hypothesis that the reductions $M_1[N/x] \xrightarrow{R}^* M_1'' * \leftarrow M_1[N'/x]$ do not contain any expansions, and in particular no δ rule, at the root position, so we can apply context closure for projections and get

$$\pi_i(M_1)[N/x] \xrightarrow{R}^* \pi_i(M_1'') * \leftarrow \pi_i(M_1)[N'/x].$$

- If $M_1 \equiv x$, and the expansion rule R is δ at the root of N , then $N' \equiv \langle \pi_1(N), \pi_2(N) \rangle$ and we can close our diagram as follows

$$\begin{array}{ccc} \pi_i(x[N/x]) & & \pi_i(x[\langle \pi_1 N, \pi_2 N \rangle / x]) \\ = & & = \\ \pi_i(N) & \xleftarrow{\pi} & \pi_i\langle \pi_1 N, \pi_2 N \rangle \end{array}$$

Here, the vertical reductions are built by context closure, while the horizontal one is a π , so no expansion rule is applied at the root in the overall reduction sequence.

- $M \equiv \langle M_1, M_2 \rangle$

We find by induction hypothesis terms M_1'' and M_2'' such that $M_1[N/x] \xrightarrow{R}^* M_1'' * \leftarrow M_1[N'/x]$ and $M_2[N/x] \xrightarrow{R}^* M_2'' * \leftarrow M_2[N'/x]$. So, we can apply the context closure rule for application and get that

$$\begin{array}{ccc} (\langle M_1, M_2 \rangle)[N/x] & & (\langle M_1, M_2 \rangle)[N'/x] \\ \langle M_1[N/x], M_2[N/x] \rangle & & \langle M_1[N'/x], M_2[N'/x] \rangle \\ \begin{array}{c} * \downarrow \\ \langle M_1'', M_2'' \rangle \end{array} & = & \begin{array}{c} * \downarrow \\ \langle M_1'', M_2'' \rangle \end{array} \end{array}$$

- $M \equiv in_C^i(M_1)$

We find by induction hypothesis a term M_1'' such that $M_1[N/x] \xrightarrow{R}^* M_1'' * \leftarrow M_1[N'/x]$. so we can apply the context closure rule for in^i and get that

$$\begin{array}{ccc} in_C^i(M_1[N/x]) & & in_C^i(M_1[N'/x]) \\ = & & = \\ in_C^i(M_1)[N/x] \xrightarrow{R}^* in_C^i(M_1'') * \leftarrow & & in_C^i(M_1)[N'/x] \end{array}$$

- $M \equiv \text{Case}(P, M_1, M_2)$

We find by induction hypothesis P'' , M_1'' and M_2'' such that $P[N/x] \xRightarrow{R} P'' * \leftarrow P[N'/x]$ and $M_1[N/x] \xRightarrow{R} M_1'' * \leftarrow M_1[N'/x]$ and $M_2[N/x] \xRightarrow{R} M_2'' * \leftarrow M_2[N'/x]$. So, we can apply the context closure rule for *Case* and get that

$$\begin{array}{ccc} \text{Case}(P, M_1, M_2)[N/x] & & \text{Case}(P, M_1, M_2)[N'/x] \\ \equiv & & \equiv \\ \text{Case}(P[N/x], (M_1)[N/x], (M_2)[N/x]) & & \text{Case}(P[N'/x], (M_1)[N'/x], (M_2)[N'/x]) \\ \begin{array}{c} * \Downarrow \\ * \Downarrow \end{array} & & \begin{array}{c} * \Downarrow \\ * \Downarrow \end{array} \\ \text{Case}(P'', M_1'', M_2'') & = & \text{Case}(P'', M_1'', M_2'') \end{array}$$

- $M \equiv (\text{rec } z : A.M_1)^i$

We assume $z \neq x$ (otherwise the result trivially holds). We find by induction hypothesis a term M_1'' such that $M_1[N/x] \xRightarrow{R} M_1'' * \leftarrow M_1[N'/x]$. so we can apply the context closure rule for *rec* and get that

$$\begin{array}{ccc} (\text{rec } z : A.M_1)^i[N/x] & & (\text{rec } z : A.M_1)^i[N'/x] \\ \equiv & & \equiv \\ (\text{rec } z : A.M_1[N/x])^i \xRightarrow{*} & & (\text{rec } z : A.M_1''^i) * \leftarrow (\text{rec } z : A.M_1[N'/x])^i \end{array}$$

□

Example 4.7 Take $M = \langle xy, x \rangle$, $N = w$ and $N' = \lambda z : A.wz$. Then

$$M[N/x] = \langle wy, w \rangle \xRightarrow{*} \langle wy, \lambda z : A.wz \rangle \leftarrow \langle (\lambda z : A.wz)y, \lambda z : A.wz \rangle = M[N'/x]$$

Looking carefully through the proof of the previous Lemma 4.6, one can see that the only cases where it is needed to apply a reverse reduction are those corresponding to an expansion rule applied at the root of N and to the presence in M of some free occurrences of x in *influential* positions. So, we can also state the following

Corollary 4.8 (Reverse reductions) Let $N \xRightarrow{R} N'$. In case R is not an expansion rule applied at the root of N (an external expansion rule) or x does not occur at an influential position in M , then $M[N/x] \xRightarrow{R} M[N'/x]$

Lemma 4.5 and 4.6 suffice to prove that all critical pairs arising from a term M by a β -reduction and another reduction rule can be solved. We can then state the following:

Proposition 4.9 (Critical Pairs are solvable)

If $M \rightarrow M'$ and $M \xRightarrow{*} M''$, then $\exists R$ such that $M' \xRightarrow{*} R$ and $M'' \xRightarrow{*} R$.

Proof. We consider every possible case of reduction from M to M' .

1. $M \xrightarrow{\beta} M'$. Thus $M \equiv (\lambda x.P)N$.

1.1. If $M \xRightarrow{*} M''$ is internal, there are two cases:

1.1.1. $P \xRightarrow{*} P'$

$$\begin{array}{ccc} (\lambda x.P)N & \xRightarrow{*} & (\lambda x.P')N \\ \beta \downarrow & & \beta \downarrow \\ P[N/x] & & P'[N/x] \end{array}$$

By lemma 4.5 we have $P[N/x] \xRightarrow{*} P'[N/x]$ or $P'[N/x] \xRightarrow{*} P[N/x]$.

1.1.2. $N \Longrightarrow N'$

$$\begin{array}{ccc} (\lambda x.P)N & \Longrightarrow & (\lambda x.P)N' \\ \beta \downarrow & & \beta \downarrow \\ P[N/x] & & P[N'/x] \end{array}$$

By lemma 4.6 there is a term R such that $P[N/x] \Longrightarrow^* R$ and $P[N'/x] \Longrightarrow^* R$.

1.2. If $M \Longrightarrow M''$ is external:

1.2.1. $M \equiv (\lambda x.P)N \xrightarrow{\eta} \lambda y.((\lambda x.P)N)y \equiv M''$

1.2.1.1. If $P[N/x]$ is not a λ -abstraction:

$$\begin{array}{ccc} (\lambda x.P)N & \xrightarrow{\eta} & \lambda y.((\lambda x.P)N)y \\ \beta \downarrow & & \beta \downarrow \\ P[N/x] & \xrightarrow{\eta} & \lambda y.P[N/x]y \end{array}$$

1.2.1.2. If $P[N/x]$ is a λ -abstraction we have two cases:

1.2.1.2.1. If P is a λ -abstraction:

$$\begin{array}{ccc} (\lambda x.(\lambda z.P'))N & \xrightarrow{\eta} & \lambda y.((\lambda x.(\lambda z.P'))N)y \\ \beta \downarrow & & \beta \downarrow \\ (\lambda z.P')[N/x] & & \lambda y.((\lambda z.P')[N/x])y \\ & & = \\ & & \lambda y.(\lambda z.(P'[N/x])y) \\ & & \beta \downarrow \\ (\lambda z.(P'[N/x])) & = & \lambda y.P'[N/x][y/z] \end{array}$$

1.2.1.2.2. If $P = x$ and N is a λ -abstraction $\lambda z.N'$:

$$\begin{array}{ccc} (\lambda x.x)\lambda z.N' & \xrightarrow{\eta} & (\lambda y.((\lambda x.x)\lambda z.N')y) \\ \beta \downarrow & & \beta \downarrow \\ \lambda z.N' & = & \lambda y.(\lambda z.N')y \\ & & \beta \downarrow \\ & & \lambda y.N'[y/z] \end{array}$$

1.2.2. $M \equiv (\lambda x.P)N \xrightarrow{\delta} \langle \pi_1((\lambda x.P)N), \pi_2((\lambda x.P)N) \rangle \equiv M''$

1.2.2.1. If $P[N/x]$ is not a pair we have:

$$\begin{array}{ccc} (\lambda x.P)N & \xrightarrow{\delta} & \langle \pi_1((\lambda x.P)N), \pi_2((\lambda x.P)N) \rangle \\ \beta \downarrow & & \beta \downarrow \\ P[N/x] & \xrightarrow{\delta} & \langle \pi_1(P[N/x]), \pi_2((\lambda x.P)N) \rangle \\ & & \beta \downarrow \\ & & \langle \pi_1(P[N/x]), \pi_2(P[N/x]) \rangle \end{array}$$

1.2.2.2. If $P[N/x]$ is a pair we have two more cases:

1.2.2.2.1. P is also a pair $\langle P_1, P_2 \rangle$:

$$\begin{array}{ccc}
(\lambda x. \langle P_1, P_2 \rangle)N & \xrightarrow{\delta} & \langle \pi_1((\lambda x. \langle P_1, P_2 \rangle)N), \pi_2((\lambda x. \langle P_1, P_2 \rangle)N) \rangle \\
\beta \downarrow & & \beta \downarrow \\
& & \langle \pi_1(\langle P_1, P_2 \rangle[N/x]), \pi_2((\lambda x. \langle P_1, P_2 \rangle)N) \rangle \\
& & \beta \downarrow \\
& & \langle \pi_1(\langle P_1, P_2 \rangle[N/x]), \pi_2(\langle P_1, P_2 \rangle[N/x]) \rangle \\
& & \pi_1 \downarrow \\
\langle P_1[N/x], P_2[N/x] \rangle & \xleftarrow{\pi_2} & \langle P_1[N/x], \pi_2(\langle P_1, P_2 \rangle[N/x]) \rangle
\end{array}$$

1.2.2.2.2. $P = x$ and N is a pair $\langle N_1, N_2 \rangle$:

$$\begin{array}{ccc}
(\lambda x. x) \langle N_1, N_2 \rangle & \xrightarrow{\delta} & \langle \pi_1((\lambda x. x) \langle N_1, N_2 \rangle), \pi_2((\lambda x. x) \langle N_1, N_2 \rangle) \rangle \\
\beta \downarrow & & \beta \downarrow \\
& & \langle \pi_1(\langle N_1, N_2 \rangle), \pi_2((\lambda x. x) \langle N_1, N_2 \rangle) \rangle \\
& & \beta \downarrow \\
& & \langle \pi_1(\langle N_1, N_2 \rangle), \pi_2(\langle N_1, N_2 \rangle) \rangle \\
& & \pi_1 \downarrow \\
\langle N_1, N_2 \rangle & \xleftarrow{\pi_2} & \langle N_1, \pi_2(\langle N_1, N_2 \rangle) \rangle
\end{array}$$

1.2.3. $M \equiv (\lambda x. P)N \xrightarrow{Top} * \equiv M''$.

Then $(\lambda x. P)N$ is of type \mathbf{T} , so also $P[N/x]$ is of type \mathbf{T} and then $P[N/x] \xrightarrow{\mathbf{T}} *$.

2. $M \xrightarrow{n} M'$.

2.1. If $M \implies M''$ is internal, then the same reduction can be performed on $\lambda z. Mz$, and the outermost term constructor of M and M'' does not change, so an expansion is still possible on M'' , and we can generally close the diagram as follows:

$$\begin{array}{ccc}
M & \implies & M'' \\
\eta \downarrow & & \eta \downarrow \\
\lambda z. Mz & \implies & \lambda z. (M'')z
\end{array}$$

2.2. If $M \implies M''$ is external, the cases to consider are:

2.2.1. $M \xrightarrow{\beta} M''$. This is the same as case 1.2.1

2.2.2. $M \xrightarrow{\pi_i} M''$. Then $M \equiv \pi_i(\langle M_1, M_2 \rangle)$ and there are two cases:

2.2.2.1. If M_i is not a λ -abstraction, the diagram looks like:

$$\begin{array}{ccc}
\pi_i(\langle M_1, M_2 \rangle) & \xrightarrow{\pi_i} & M_i \\
\eta \downarrow & & \eta \downarrow \\
\lambda z. \pi_i(\langle M_1, M_2 \rangle)z & \xrightarrow{\pi_i} & \lambda z. M_i z
\end{array}$$

2.2.2.2. If M_i is a λ -abstraction $\lambda y.M'_i$, the diagram looks like:

$$\begin{array}{ccc}
\pi_i(\langle M_1, M_2 \rangle) & \xrightarrow{\pi_i} & \lambda y.M'_i \\
\downarrow \eta & & \\
\lambda z.\pi_i(\langle M_3, M_2 \rangle)z & & \\
\pi_i \downarrow & & \\
\lambda z.(\lambda y.M'_i)z & & \\
\beta \downarrow & & \\
\lambda z.(M'_i[z/y]) & = &
\end{array}$$

2.2.3. $M \xrightarrow{\rho} M''$. Then $M \equiv \text{Case}(in_{C_1+C_2}^i(P), M_1, M_2)$

$$\begin{array}{ccc}
\text{Case}(in_{C_1+C_2}^i(P), M_1, M_2) & \xrightarrow{\rho} & M_i P \\
\eta \downarrow & & \eta \downarrow \\
\lambda z.\text{Case}(in_{C_1+C_2}^i(P), M_1, M_2)z & \xrightarrow{\rho} & \lambda z.(M_i P)z
\end{array}$$

2.2.4. $M \xrightarrow{rec} M''$. Then $M = (rec\ y.M_1)^i$ and there are two possible cases:

2.2.4.1. If M_1 is not a λ -abstraction:

$$\begin{array}{ccc}
(rec\ y.M_1)^i & \xrightarrow{rec} & M_1[(rec\ y.M_1)^{i-1}/y] \\
\eta \downarrow & & \eta \downarrow \\
\lambda z.(rec\ y.M_1)^i z & \xrightarrow{rec} & \lambda z.(M_1[(rec\ y.M_1)^{i-1}/y])z
\end{array}$$

2.2.4.2. If $M_1 \equiv \lambda w.M'_1$:

$$\begin{array}{ccc}
(rec\ y.(\lambda w.M'_1))^i & \xrightarrow{rec} & (\lambda w.M'_1)[(rec\ y.(\lambda w.M'_1))^{i-1}/y] \\
\eta \downarrow & & \\
\lambda z.(rec\ y.\lambda w.M'_1)^i z & & \\
rec \downarrow & & \\
\lambda z.(\lambda w.M'_1)[(rec\ y.(\lambda w.M'_1))^{i-1}/y]z & & \\
\beta \downarrow & & \\
\lambda z.(M'_1[(rec\ y.(\lambda w.M'_1))^{i-1}/y][z/w]) & = &
\end{array}$$

3. $M \xrightarrow{Top} *$. Since M is of type \mathbf{T} and $M \Longrightarrow M''$, also M'' is of type \mathbf{T} by proposition 3.7. Then $M'' \xrightarrow{Top} *$.

4. $M \xrightarrow{\pi_i} M'$.

4.1. If $M \Longrightarrow M''$ is internal:

4.1.1. $M \equiv \pi_1(\langle M_1, M_2 \rangle) \Longrightarrow \pi_1(\langle M'_1, M_2 \rangle) \equiv M'$, where $M_1 \Longrightarrow M'_1$

$$\begin{array}{ccc}
\pi_1(\langle M_1, M_2 \rangle) & \Longrightarrow & \pi_1(\langle M'_1, M_2 \rangle) \\
\pi_1 \downarrow & & \pi_1 \downarrow \\
M_1 & \Longrightarrow & M'_1
\end{array}$$

- Idem for $M \equiv \pi_2(\langle M_1, M_2 \rangle) \implies \pi_2(\langle M_1, M'_2 \rangle) \equiv M'$, where $M_2 \implies M'_2$.
- 4.1.2. $M \equiv \pi_1(\langle M_1, M_2 \rangle) \implies \pi_1(\langle M_1, M'_2 \rangle) \equiv M'$, where $M_2 \implies M'_2$.

$$\begin{array}{ccc} \pi_1(\langle M_1, M_2 \rangle) & \implies & \pi_1(\langle M_1, M'_2 \rangle) \\ \pi_1 \downarrow & & \pi_1 \downarrow \\ M_1 & \equiv & M_1 \end{array}$$

Idem for $M \equiv \pi_2(\langle M_1, M_2 \rangle) \implies \pi_2(\langle M'_1, M_2 \rangle) \equiv M'$, where $M_1 \implies M'_1$.

4.2. If $M \implies M''$ is external:

- 4.2.1. $M \xrightarrow{\eta} M''$. This is the same as case 2.2.2.
- 4.2.2. $M \xrightarrow{Top} M''$. This is considered in case 3.
- 4.2.3. $M \xrightarrow{\delta} M''$. Then $M \equiv \pi_i(\langle M_1, M_2 \rangle)$.

4.2.3.1. If M_i is not a pair:

$$\begin{array}{ccc} \pi_i(\langle M_1, M_2 \rangle) & \xrightarrow{\delta} & \langle \pi_1(\pi_i(\langle M_1, M_2 \rangle)), \pi_2(\pi_i(\langle M_1, M_2 \rangle)) \rangle \\ \pi_i \downarrow & & \pi_i \downarrow \\ & & \langle \pi_1(M_i), \pi_2(\pi_i(\langle M_1, M_2 \rangle)) \rangle \\ & & \pi_i \downarrow \\ M_i & \xrightarrow{\delta} & \langle \pi_1(M_i), \pi_2(M_i) \rangle \end{array}$$

4.2.3.2. If M_i is a pair $\langle P_1, P_2 \rangle$:

$$\begin{array}{ccc} \pi_i(\langle M_1, M_2 \rangle) & \xrightarrow{\delta} & \langle \pi_1(\pi_i(\langle M_1, M_2 \rangle)), \pi_2(\pi_i(\langle M_1, M_2 \rangle)) \rangle \\ \pi_i \downarrow & & \pi_i \downarrow \\ & & \langle \pi_1(\langle P_1, P_2 \rangle), \pi_2(\pi_1(\langle M_1, M_2 \rangle)) \rangle \\ & & \pi_i \downarrow \\ & & \langle \pi_1(\langle P_1, P_2 \rangle), \pi_2(\langle P_1, P_2 \rangle) \rangle \\ & & \pi_1 \downarrow \\ \langle P_1, P_2 \rangle & \xleftarrow{\pi_2} & \langle P_1, \pi_2(\langle P_1, P_2 \rangle) \rangle \end{array}$$

5. $M \xrightarrow{\delta} M'$.

5.1. If $M \implies M''$ is internal, then the same reduction can be performed on $\langle \pi_1(M), \pi_2(M) \rangle$, and the outermost term constructor of M and M'' does not change, so an expansion is still possible on M'' , and we can generally close the diagram as follows:

$$\begin{array}{ccc} M & \implies & M'' \\ \delta \downarrow & & \delta \downarrow \\ \langle \pi_1(M), \pi_2(M) \rangle & \implies & \langle \pi_1(M), \pi_2(M'') \rangle \end{array}$$

5.2. If $M \implies M''$ is external:

- 5.2.1. $M \xrightarrow{\beta} M''$. This is the same as case 1.2.2
- 5.2.2. $M \xrightarrow{\pi_i} M''$. This is the same as case 4.2.3

5.2.3. $M \equiv \text{Case}(in_{C_1+C_2}^i(P), M_1, M_2) \xrightarrow{\rho} M_i P \equiv M''$.

$$\begin{array}{ccc} \text{Case}(in_{C_1+C_2}^i(P), M_1, M_2) & \xrightarrow{\rho} & M_i P \\ \delta \downarrow & & \delta \downarrow \\ \langle \pi_1(M), \pi_2(M) \rangle & \xrightarrow{\rho^*} & \langle \pi_1(M_i P), \pi_2(M_i P) \rangle \end{array}$$

5.2.4. $M \equiv (\text{rec } y : C.P)^i \rightarrow P[(\text{rec } y : C.P)^{i-1}/y] \equiv M''$.

5.2.4.1. If P is not a pair:

$$\begin{array}{ccc} M \equiv (\text{rec } y : C.P)^i & \xrightarrow{\quad} & P[(\text{rec } y : C.P)^{i-1}/y] \\ \delta \downarrow & & \delta \downarrow \\ \langle \pi_1(M), \pi_2(M) \rangle & \xrightarrow{\text{rec } *} & \langle \pi_1(P[M/y]), \pi_2(P[M/y]) \rangle \end{array}$$

5.2.4.2. If P is a pair $\langle P_1, P_2 \rangle$:

$$\begin{array}{ccc} M \equiv (\text{rec } y : C.\langle P_1, P_2 \rangle)^i & \xrightarrow{\quad} & \langle P_1, P_2 \rangle[M/y] \\ \delta \downarrow & & \\ \langle \pi_1(M), \pi_2(M) \rangle & & \\ \downarrow \text{rec} & & \\ \langle \pi_1(\langle P_1, P_2 \rangle[M/y]), \pi_2(M) \rangle & & \\ \downarrow \text{rec} & & \\ \langle \pi_1(\langle P_1, P_2 \rangle[M/y]), \pi_2(\langle P_1, P_2 \rangle[M/y]) \rangle & & \\ \downarrow \pi_1 & & \\ \langle P_1[M/y], \pi_2(\langle P_1, P_2 \rangle[M/y]) \rangle & & \\ \downarrow \pi_2 & & \\ \langle P_1[M/y], P_2[M/y] \rangle & = & \end{array}$$

6. $M \xrightarrow{\rho} M'$.

6.1. If $M \Rightarrow M''$ is internal:

6.1.1. $\text{Case}(in_C^i(P), M_1, M_2) \Rightarrow \text{Case}(in_C^i(P'), M_1, M_2)$, where $P \Rightarrow P'$.

$$\begin{array}{ccc} \text{Case}(in_C^i(P), M_1, M_2) & \Rightarrow & \text{Case}(in_C^i(P'), M_1, M_2) \\ \rho \downarrow & & \rho \downarrow \\ M_i P & \xrightarrow{\quad} & M_i P' \end{array}$$

6.1.2. $\text{Case}(in_C^1(P), M_1, M_2) \Rightarrow \text{Case}(in_C^1(P), M'_1, M_2)$, where $M_1 \Rightarrow M'_1$ but $M_1 \xrightarrow{\neg n} M'_1$

$$\begin{array}{ccc} \text{Case}(in_C^1(P), M_1, M_2) & \Rightarrow & \text{Case}(in_C^1(P), M'_1, M_2) \\ \rho \downarrow & & \rho \downarrow \\ M_1 P & \xrightarrow{\quad} & M'_1 P \end{array}$$

Idem for $Case(in_C^2(P), M_1, M_2) \Longrightarrow Case(in_C^2(P), M_1, M'_2)$, where $M_2 \xrightarrow{\eta} M'_2$.
6.1.3. $Case(in_C^1(P), M_1, M_2) \Longrightarrow Case(in_C^1(P), M'_1, M_2)$, where $M_1 \longrightarrow \lambda x.M_1x$

$$Case(in_C^1(P), M_1, M_2) \Longrightarrow Case(in_C^1(P), \lambda x.M_1x, M_2)$$

$$\begin{array}{ccc} \rho \downarrow & & \rho \downarrow \\ M_1P & \xleftarrow{\beta} & (\lambda x.M_1x)P \end{array}$$

Idem for $Case(in_C^2(P), M_1, M_2) \Longrightarrow Case(in_C^2(P), M_1, M'_2)$, where $M_2 \longrightarrow \lambda x.M_2x$

6.2. If $M \Longrightarrow M''$ is external:

- 6.2.1. $M \xrightarrow{\eta} M''$. This is the same as case 2.2.3.
- 6.2.2. $M \xrightarrow{Top} M''$. This is considered in case 3.
- 6.2.3. $M \xrightarrow{\delta} M''$. This is the same as case 5.2.3.

7. $M \xrightarrow{rec} M'$.

7.1. If $M \Longrightarrow M''$ is internal:

- 7.1.1. $(rec\ y : C.P)^i \Longrightarrow (rec\ y : C.P')^i$, where $P \Longrightarrow P'$. We have $P[(rec\ y : C.P)^{i-1}/y] \Longrightarrow P[(rec\ y : C.P')^{i-1}/y]$ by lemma 4.8 and by lemma 4.5 $P[(rec\ y : C.P')^{i-1}/y] \Longrightarrow^* P'[(rec\ y : C.P')^{i-1}/y]$ or $P'[(rec\ y : C.P')^{i-1}/y] \Longrightarrow^* P[(rec\ y : C.P')^{i-1}/y]$.
In the first case:

$$\begin{array}{ccc} (rec\ y : C.P)^i & \Longrightarrow & (rec\ y : C.P')^i \\ \text{rec} \downarrow & & \text{rec} \downarrow \\ P[(rec\ y : C.P)^{i-1}/y] & & P[(rec\ y : C.P')^{i-1}/y] \\ * \Downarrow & & \downarrow \\ P[(rec\ y : C.P')^{i-1}/y] & \Longrightarrow^* & P'[(rec\ y : C.P')^{i-1}/y] \end{array}$$

In the second case:

$$\begin{array}{ccc} (rec\ y : C.P)^i & \Longrightarrow & (rec\ y : C.P')^i \\ \text{rec} \downarrow & & \text{rec} \downarrow \\ P[(rec\ y : C.P)^{i-1}/y] & & P[(rec\ y : C.P')^{i-1}/y] \\ * \Downarrow & & \downarrow \\ P[(rec\ y : C.P')^{i-1}/y] & \Longleftarrow^* & P'[(rec\ y : C.P')^{i-1}/y] \end{array}$$

7.2. If $M \Longrightarrow M''$ is external:

- 7.2.1. $M \xrightarrow{\eta} M''$. This is the same as case 2.2.4.
- 7.2.2. $M \xrightarrow{Top} M''$. This is considered in case 3.
- 7.2.3. $M \xrightarrow{\delta} M''$. This is the same as case 5.2.4.

□

4.3 From Solved Critical Pairs to Full Weak Confluence

It is to be noted that the solvability of critical pairs we just proved as Proposition 4.9 does not allow us to deduce the weak confluence of the calculus via the famous Knuth-Bendix Critical Pairs Lemma. That Lemma holds only for algebraic rewrite systems, and not for the λ -calculus, that has the higher order rewrite rule β . We need to prove local confluence explicitly, and to do so the following remark is useful.

Remark 4.10 (Expansion rules) *In case the two reductions $M' \leftarrow M \Rightarrow M''$ do not involve η (resp. δ) rules applied at the root positions of M , it is possible to close the diagram without using η (resp. δ) rules at the root, except in the three cases shown below: external π 's and internal η , external β and internal δ . Notice that M is not a λ - abstraction in the first diagram, N is not a λ - abstraction in the second and $M[N/x]$ is not a pair in the third one.*

$$\begin{array}{ccc}
\pi_1(\langle M, N \rangle) \xRightarrow{\eta} \pi_1(\langle \lambda x.Mx, N \rangle) & \pi_2(\langle M, N \rangle) \xRightarrow{\eta} \pi_2(\langle M, \lambda x.Nx \rangle) \\
\pi \downarrow & \Downarrow \pi & \pi \downarrow & \Downarrow \pi \\
M \xRightarrow{\eta} \lambda x.Mx & & N \xRightarrow{\eta} \lambda x.Nx \\
\\
(\lambda x : A.M)N \xRightarrow{\delta} (\lambda x : A.\langle \pi_1(M), \pi_2(M) \rangle)N \\
\beta \downarrow & & \Downarrow \beta \\
M[N/x] \xRightarrow{\delta} \langle \pi_1(M[N/x]), \pi_2(M[N/x]) \rangle
\end{array}$$

With this additional knowledge, we can prove that \Rightarrow is actually weakly confluent.

Theorem 4.11 (Weak Confluence) *If $M' \leftarrow M \Rightarrow M''$ then there exist a term M''' such that $M' \Rightarrow^* M''' \ast \leftarrow M''$ (i.e. the reduction relation \Rightarrow is weakly confluent). Furthermore, if the reductions in $M' \leftarrow M \Rightarrow M''$ do not contain η (resp. δ) rules applied at the root of M , it is possible also to close the diagram without applying η (resp. δ) rules at the root, except in the cases shown in the previous Remark 4.10.*

Proof. We will prove that there exists a term M''' such that $M' \Rightarrow^* M''' \ast \leftarrow M''$, by induction on the derivation of $M \Rightarrow M'$. First of all, we remark that if one of the two one-step reductions $M \Rightarrow M'$ and $M \Rightarrow M''$ is actually an external reduction $M \xrightarrow{M} M'$ and $M \xrightarrow{M} M''$, then the result comes directly from Proposition 4.9. So we will need to consider in the following only the cases where both reductions are internal reductions.

We proceed now by cases on the last rule used to derive $M \Rightarrow M'$.

- $M \equiv (M_1 M_2) \Rightarrow (M'_1 M_2) \equiv M'$ comes from $M_1 \Rightarrow M'_1$. In this case, the η rule cannot be applied at the root position of M_1 because M_1 is evaluated. Then we have two cases:

- the reduction $M \equiv (M_1 M_2) \Rightarrow (M''_1 M_2) \equiv M''$ comes from a reduction $M_1 \Rightarrow M''_1$. Now we have to consider two cases:

- * $M'_1 \leftarrow M_1 \Rightarrow M''_1$ is not one of the exceptional cases for η of the Remark 4.10: then we know that there are no η at the root position in $M'_1 \Rightarrow^* M'''_1 \ast \leftarrow M''_1$. By induction hypothesis we get a term M'''_1 that can be used to close the diagram $M'_1 \leftarrow M_1 \Rightarrow M''_1$ via $M'_1 \Rightarrow^* M'''_1 \ast \leftarrow M''_1$, and we can close our original diagram with

$$M' \equiv (M'_1 M_2) \Rightarrow^* (M'''_1 M_2) \ast \leftarrow (M''_1 M_2) \equiv M''$$

- * $M'_1 \leftarrow M_1 \Rightarrow M''_1$ is one of the exceptional cases for η , hence M_1 is $\pi_1(\langle P, Q \rangle)$ for some terms P and Q . We can still close the original diagram as follows:

$$\begin{array}{ccccccc}
(\pi_1(\langle P, Q \rangle))M_2 \xRightarrow{\eta} (\pi_1(\langle \lambda x.Px, Q \rangle))M_2 & & (\pi_2(\langle P, Q \rangle))M_2 \xRightarrow{\eta} (\pi_2(\langle P, \lambda x.Qx \rangle))M_2 \\
\Downarrow \pi & & \Downarrow \pi & & \Downarrow \pi & & \Downarrow \pi \\
PM_2 & \equiv & (\lambda x.Px)M_2 & & QM_2 & \equiv & (\lambda x.Qx)M_2 \\
\Downarrow \beta & & \Downarrow \beta & & \Downarrow \beta & & \Downarrow \beta \\
PM_2 & & PM_2 & & QM_2 & & QM_2
\end{array}$$

- the reduction $M \equiv (M_1 M_2) \Longrightarrow (M_1 M_2'') \equiv M''$ comes from a reduction $M_2 \Longrightarrow M_2''$. We can close the diagram using the same original reductions,

$$M' \equiv (M_1' M_2) \Longrightarrow (M_1' M_2'') \Leftarrow (M_1 M_2'') \equiv M''$$

because we know that η is not applied to M_1 to get to M_1' .

- $M \equiv (M_1 M_2) \Longrightarrow (M_1 M_2') \equiv M'$ comes from $M_2 \Longrightarrow M_2'$. Then we have two cases:
 - the reduction $M \equiv (M_1 M_2) \Longrightarrow (M_1 M_2'') \equiv M''$ comes from a reduction $M_2 \Longrightarrow M_2''$. By induction hypothesis we get a term M_2''' that can be used to close $M_2' \Leftarrow M_2 \Longrightarrow M_2''$ via $M_2' \Longrightarrow^* M_2''' \Leftarrow^* M_2''$. Now $M' \equiv (M_1 M_2') \Longrightarrow^* (M_1 M_2'') \Leftarrow^* (M_1 M_2'') \equiv M''$ can be used to close our original diagram.
 - the reduction $M \equiv (M_1 M_2) \Longrightarrow (M_1'' M_2) \equiv M''$ comes from a reduction $M_1 \Longrightarrow M_1''$. In this case, we know that η cannot be applied at the top to M_1 to get to M_1'' because M_1 is evaluated. So, we can close the diagram using the same original reductions, $M' \equiv (M_1 M_2') \Longrightarrow (M_1'' M_2') \Leftarrow (M_1'' M_2') \equiv M''$.
- $M \equiv (\pi_i(M_1)) \Longrightarrow (\pi_i(M_1')) \equiv M'$ comes from $M_1 \Longrightarrow M_1'$ and $M \equiv (\pi_i(M_1)) \Longrightarrow (\pi_i(M_1'')) \equiv M''$ comes from $M_1 \Longrightarrow M_1''$. Then neither $M_1 \Longrightarrow M_1'$ nor $M_1 \Longrightarrow M_1''$ can use δ rules at the root of M because it is projected.

Now we have two cases:

- $M_1 \Longrightarrow M_1'$ and $M_1 \Longrightarrow M_1''$ are not the exceptional cases for δ of Remark 4.10. By induction hypothesis there is an M_1''' s.t. $M_1' \Longrightarrow^* M_1''' \Leftarrow^* M_1''$ without δ rules at the root, and we can close our diagram by $\pi_i(M_1') \Longrightarrow^* \pi_i(M_1''') \Leftarrow^* \pi_i(M_1'')$.
- $M_1 \Longrightarrow M_1'$ and $M_1 \Longrightarrow M_1''$ is the exceptional case for δ , so $M_1 \equiv (\lambda x.P)Q$ for some terms P and Q . We can still close our original diagram as follows:

$$\begin{array}{ccc}
\pi_i((\lambda x.P)Q) & \xrightarrow{\delta} & \pi_i((\lambda x.\langle \pi_1(P), \pi_2(P) \rangle)Q) \\
\downarrow \beta & & \downarrow \beta \\
& & \pi_i(\langle \pi_1(P[Q/x]), \pi_2(P[Q/x]) \rangle) \\
& & \downarrow \pi \\
\pi_i(P[Q/x]) & \xrightarrow{\delta} & \pi_i(P[Q/x])
\end{array}$$

- $M \equiv \lambda x.M_1 \Longrightarrow \lambda x.M_1' \equiv M'$ comes from $M_1 \Longrightarrow M_1'$ and $M \equiv \lambda x.M_1 \Longrightarrow \lambda x.M_1'' \equiv M''$ comes from $M_1 \Longrightarrow M_1''$. By induction hypothesis there is an M_1''' s.t. $M_1' \Longrightarrow^* M_1''' \Leftarrow^* M_1''$ and we can close our diagram by $\lambda x.M_1' \Longrightarrow^* \lambda x.M_1''' \Leftarrow^* \lambda x.M_1''$.
- $M \equiv \langle M_1, M_2 \rangle \Longrightarrow \langle M_1', M_2 \rangle \equiv M'$ comes from $M_1 \Longrightarrow M_1'$. Now we have to consider two cases:

- the reduction $M \equiv \langle M_1, M_2 \rangle \Longrightarrow \langle M_1, M_2'' \rangle \equiv M''$ comes from a reduction $M_2 \Longrightarrow M_2''$. By induction hypothesis there is a term M_1''' s.t. we can close the diagram $M_1' \Leftarrow M_1 \Longrightarrow M_1''$ via $M_1' \Longrightarrow^* M_1''' \Leftarrow^* M_1''$, and we can close our original diagram with

$$M' \equiv \langle M_1', M_2 \rangle \Longrightarrow^* \langle M_1''', M_2 \rangle \Leftarrow^* \langle M_1'', M_2 \rangle \equiv M''$$

- the reduction $M \equiv \langle M_1, M_2 \rangle \Longrightarrow \langle M_1, M_2'' \rangle \equiv M''$ comes from a reduction $M_2 \Longrightarrow M_2''$. We can close the diagram using the same original reductions,

$$M' \equiv \langle M_1, M_2 \rangle \Longrightarrow \langle M_1, M_2'' \rangle \Leftarrow \langle M_1, M_2'' \rangle \equiv M''$$

- $M \equiv \text{in}_C^i(M_1) \Longrightarrow \text{in}_C^i(M'_1) \equiv M'$ comes from $M_1 \Longrightarrow M'_1$ and $M \equiv \text{in}_C^i(M_1) \Longrightarrow \text{in}_C^i(M''_1) \equiv M''$ comes from $M_1 \Longrightarrow M''_1$. By induction hypothesis there is an M'''_1 s.t. $M'_1 \Longrightarrow^* M'''_1 \ast \Leftarrow M''_1$ and we can close our diagram by $\text{in}_C^i(M'_1) \Longrightarrow^* \text{in}_C^i(M'''_1) \ast \Leftarrow \text{in}_C^i(M''_1)$.
- $M \equiv \text{rec } x : A.M_1 \Longrightarrow \text{rec } x : A.M'_1 \equiv M'$ comes from $M_1 \Longrightarrow M'_1$ and $M \equiv \text{rec } x : A.M_1 \Longrightarrow \text{rec } x : A.M''_1 \equiv M''$ comes from $M_1 \Longrightarrow M''_1$. Then we can find by induction hypothesis an M'''_1 s.t. $M'_1 \Longrightarrow^* M'''_1 \ast \Leftarrow M''_1$ and we can close our diagram by $\text{rec } x : A.M'_1 \Longrightarrow^* \text{rec } x : A.M'''_1 \ast \Leftarrow \text{rec } x : A.M''_1$.
- We are left to consider the case of $M \equiv \text{Case}(P, M_1, M_2)$.
 - To avoid a mechanical repetition of similar proofs, notice that if the internal reduction to M' and M'' are performed on different subterms, then we can close the diagram by commuting the two reductions. We show just one case.

$$\begin{array}{ccc}
\text{Case}(P, M_1, M_2) & \xRightarrow{R2} & \text{Case}(P, M'_1, M_2) \\
R1 \Downarrow & & \Downarrow R1 \\
\text{Case}(P', M_1, M_2) & \xRightarrow{R2} & \text{Case}(P', M'_1, M_2)
\end{array}$$

- If the internal reduction to M' and M'' are performed on the same subterm Q , say $Q' \xleftarrow{R1} Q \xRightarrow{R2} Q''$, then there is a Q''' , by induction hypothesis, s.t. $Q' \Longrightarrow^* Q''' \ast \Leftarrow Q''$, and we can close the diagram by extending these last reductions to the *Case* expression. Again, we detail just one case.

$$\begin{array}{ccc}
\text{Case}(P, M_1, M_2) & \xRightarrow{R2} & \text{Case}(P'', M_1, M_2) \\
R1 \Downarrow & & \Downarrow \\
\text{Case}(P', M_1, M_2) & \Longrightarrow & \text{Case}(P''', M'_1, M_2)
\end{array}$$

□

5 Strong Normalization

We provide in this section the proof of strong normalization for our calculus. The key idea is to reduce strong normalization of the system with expansion rules to that of the system without expansion rules and for this, we show how the calculus without expansions can be used to simulate the calculus with expansions. We will use a fundamental property relating strong normalization of two systems:

Proposition 5.1 *Let \mathcal{R}_1 and \mathcal{R}_2 be two reduction systems and \mathcal{T} a translation from terms in \mathcal{R}_1 to terms in \mathcal{R}_2 . If for every reduction $M_1 \xRightarrow{\mathcal{R}_1} M_2$ there is a non empty reduction sequence $P_1 \xRightarrow{\mathcal{R}_2}^+ P_2$ such that $\mathcal{T}(M_i) = P_i$, for $i = 1, 2$, then the strong normalization of \mathcal{R}_2 implies that of \mathcal{R}_1 .*

Proof. Suppose \mathcal{R}_2 is strongly normalizing and \mathcal{R}_1 is not. Then there is an infinite reduction sequence $M_1 \xRightarrow{\mathcal{R}_1} M_2 \xRightarrow{\mathcal{R}_1} \dots$ and from this reduction we can construct an infinite reduction sequence $\mathcal{T}(M_1) \xRightarrow{\mathcal{R}_2}^+ \mathcal{T}(M_2) \xRightarrow{\mathcal{R}_2}^+ \dots$ which leads to a contradiction. □

The goal is now to find a translation of terms mapping our calculus into itself such that for every possible reduction in the original system from a term M to another term N , there is a reduction sequence from the translation of M to the translation of N , that is *non empty* and *does not* contain any expansion. Then the previous proposition allows us to derive the strong normalization property for the full system from that of the system without expansion rules, which can be proved using standard techniques.

5.1 Simulating Expansions without Expansions

The first naïve idea that comes to the mind is to choose a translation such that expansion rules are completely impossible on a translated term. This essentially amounts to associate to a term M its η - δ normal form, so that translating a term corresponds then to executing all the possible expansions.

Unfortunately, this simple solution is not a good one: if M reduces to N via an expansion, then the translation of M and that of N are the same term, so to such a reduction step in the full system corresponds an *empty* reduction sequence in the translation, and this does not allow us to apply proposition 5.1.

This leads us to consider a more sophisticated translation that maps a term M to a term M° where expansions are not fully executed as above, but just *marked* in such a way that they can be executed during the simulation process, if necessary, by a rule that is not an expansion.

Let us see how to do this on a simple example: take a variable z of type $A_1 \times A_2$, where the A_i 's are atomic types different from \mathbf{T} . By performing a δ expansion we obtain its normal form w.r.t. expansion rules: $\langle \pi_1(z), \pi_2(z) \rangle$. Instead of executing this reduction, we just *mark* it in the translation by applying to z an appropriate *expansor* term $\lambda x : A_1 \times A_2. \langle \pi_1(x), \pi_2(x) \rangle$. As for $\langle \pi_1(z), \pi_2(z) \rangle$, it is in normal form w.r.t. expansions, so the translation does not modify it in any way. Now, we have the reduction sequence

$$z^\circ \equiv (\lambda x : A_1 \times A_2. \langle \pi_1(x), \pi_2(x) \rangle)z \rightarrow_\beta \langle \pi_1(z), \pi_2(z) \rangle$$

where the translation of z reduces to the translation of $\langle \pi_1(z), \pi_2(z) \rangle$, and the δ expansion from z to $\langle \pi_1(z), \pi_2(z) \rangle$ is simulated in the translation by a β -rule. Clearly, in a generic term M there are many positions where an expansion can be performed, so the translation will have to take into account the *structure* of M and insert the appropriate expandors at all these positions⁵.

Anyway, expandors must be carefully defined to correctly represent not only the expansion step arising from a redex already present in M , but also all the expansion sequences that such step can create: if in the previous example the type A_1 is taken to be an arrow type and the type A_2 a product type, then the term $\pi_1(z)$ can be further η -expanded and the term $\pi_2(z)$ can be expanded by a δ -rule, and the expansor $\lambda x : A_1 \times A_2. \langle \pi_1(x), \pi_2(x) \rangle$ cannot simulate these further possible reductions. This can only be done by storing in the expansor terms all the information on possible future expansions, that is fully contained in the *type* of the term we are marking.

Definition 5.2 (Translation) *To every type C we associate a term, called the expansor of type C and denoted Δ_C , defined by induction as follows:*

$$\begin{aligned} \Delta_{A \rightarrow B} &= \lambda x : A \rightarrow B. \lambda z : A. \Delta_B(x(\Delta_A z)) \\ \Delta_{A \times B} &= \lambda x : A \times B. \langle \Delta_A(\pi_1(x)), \Delta_B(\pi_2(x)) \rangle \\ \Delta_A &\text{ is empty, in any other case} \end{aligned}$$

We then define a translation M° for a term $M : A$ as follows:

$$M^\circ = \begin{cases} M^{\circ\circ} & \text{if } M \text{ is a } \lambda\text{-abstraction or a pair} \\ \Delta_A^k M^{\circ\circ} & \text{for any } k > 0 \text{ otherwise} \end{cases}$$

where $\Delta_A^k M$ denotes the term $\underbrace{(\Delta_A \dots (\Delta_A M) \dots)}_{k \text{ times}}$ and $M^{\circ\circ}$ is defined by induction as:

$$\begin{array}{llll} x^{\circ\circ} & = & x & (\lambda x : B. M)^{\circ\circ} & = & \lambda x : B. M^\circ \\ *^{\circ\circ} & = & * & (\text{rec } y : A. M)^{i^{\circ\circ}} & = & (\text{rec } y : A. M^\circ)^i \\ \langle M, N \rangle^{\circ\circ} & = & \langle M^\circ, N^\circ \rangle & \text{Case}(R, M, N)^{\circ\circ} & = & \text{Case}(R^\circ, M^\circ, N^\circ) \\ (MN)^{\circ\circ} & = & (M^{\circ\circ} N^\circ) & \pi_i(M)^{\circ\circ} & = & \pi_i(M^{\circ\circ}) \\ \text{in}_C^i(M)^{\circ\circ} & = & \text{in}_C^i(M^\circ) & & & \end{array}$$

⁵Notice that we cannot insert expandors in influential positions: if a term M is expanded, say to $\langle \pi_1(M), \pi_2(M) \rangle$, then its root becomes an influential position, and we cannot insure that the translation of M reduces to a translation of $\langle \pi_1(M), \pi_2(M) \rangle$: expandors get used, and after some reduction steps we end up with a naked pair not preceded by an expansor.

This corresponds exactly to the marking procedure described before, but for a little detail: in the translation we allow *any* number of markers to be used (the integer k can be any positive number), and not just one as seemed to suffice for the examples above.

The need for this additional twist in the definition is best understood with an example. Consider two atomic types A and B and the term $(\lambda x : A \times B.x)z$: if k is fixed to be one (*i.e.* we allow only one expander as marker) then its translation $((\lambda x : A \times B.x)z)^\circ$ is $\Delta_{A \times B}((\lambda x : A \times B.\Delta_{A \times B}x)\Delta_{A \times B}z)$. Now $(\lambda x : A \times B.x)z \xrightarrow{\beta} z$, so we have to verify that $((\lambda x : A \times B.x)z)^\circ$ reduces to z° in at least one step. We have:

$$\Delta_{A \times B}((\lambda x : A \times B.\Delta_{A \times B}x)\Delta_{A \times B}z) \Longrightarrow \Delta_{A \times B}\Delta_{A \times B}\Delta_{A \times B}z$$

However, even if both $\Delta_{A \times B}^3 z$ and $\Delta_{A \times B}z$ reduce to the same term $\langle \pi_1(z), \pi_2(z) \rangle$, it is not true that $\Delta_{A \times B}^3 z \Longrightarrow^* \Delta_{A \times B}z$. Anyway, if we admit $\Delta_{A \times B}^3 z$ as a possible translation of z we will have the desired property relating reductions and translations. Hence, to be precise, our method associates to each term not just one translation, but a whole family of possible translations, all with the same structure, but with different numbers of expanders used as markers.

What is important for our proof is that when we are given a reduction $M_1 \Longrightarrow M_2 \dots \Longrightarrow M_n$ in the full calculus, then no matter which possible translation M_1° we choose for M_1 , the reductions used in the simulation process all go through possible translations M_i° of the M_i .

Translations preserve types and leave unchanged terms where expansions are not possible.

Lemma 5.3 *If $\Gamma \vdash M : A$, then $\Gamma \vdash (\Delta_A M) : A$.*

Proof. By induction on the structure of A .

- If A is neither a functional, nor a product type, then Δ_A is empty and the property trivially holds.
- $A \equiv B \rightarrow C$. Since $\Gamma, x : B \rightarrow C, z : B \vdash z : B$, we have by induction hypothesis $\Gamma, x : B \rightarrow C, z : B \vdash (\Delta_B z) : B$

$$\frac{\Gamma, x : B \rightarrow C, z : B \vdash x : B \rightarrow C \quad \Gamma, x : B \rightarrow C, z : B \vdash (\Delta_B z) : B}{\Gamma, x : B \rightarrow C, z : B \vdash (x(\Delta_B z)) : C}$$

Again by induction hypothesis $\Gamma, x : B \rightarrow C, z : B \vdash \Delta_C(x(\Delta_B z)) : C$ and thus:

$$\frac{\frac{\frac{\Gamma, x : B \rightarrow C, z : B \vdash \Delta_C(x(\Delta_B z)) : C}{\Gamma, x : B \rightarrow C \vdash \lambda z : B.\Delta_C(x(\Delta_B z)) : B \rightarrow C} \quad \Gamma \vdash M : B \rightarrow C}{\Gamma \vdash \lambda x : B \rightarrow C.\lambda z : B.\Delta_C(x(\Delta_B z)) : (B \rightarrow C) \rightarrow (B \rightarrow C)}}{\Gamma \vdash (\Delta_{B \rightarrow C} M) : B \rightarrow C}$$

- $A \equiv B \times C$. Since $\Gamma, x : B \times C \vdash x : B \times C$, then $\Gamma, x : B \times C \vdash \pi_1(x) : B$ and $\Gamma, x : B \times C \vdash \pi_2(x) : C$. By induction hypothesis $\Gamma, x : B \times C \vdash \Delta_B \pi_1(x) : B$ and $\Gamma, x : B \times C \vdash \Delta_C \pi_2(x) : C$.

$$\frac{\frac{\frac{\Gamma, x : B \times C \vdash \Delta_B \pi_1(x) : B \quad \Gamma, x : B \times C \vdash \Delta_C \pi_2(x) : C}{\Gamma, x : B \times C \vdash \langle \Delta_B \pi_1(x), \Delta_C \pi_2(x) \rangle : B \times C} \quad \Gamma \vdash M : B \times C}{\Gamma \vdash \lambda x : B \times C.\langle \Delta_B \pi_1(x), \Delta_C \pi_2(x) \rangle : (B \times C) \rightarrow (B \times C)}}{\Gamma \vdash \Delta_{B \times C} M : B \times C}$$

□

Corollary 5.4 *If $\Gamma \vdash M : A$, then $\Gamma \vdash \Delta_A^k M : A$, for any $k \geq 0$.*

Lemma 5.5 (Type Preservation) *If $\Gamma \vdash M : A$, then $\Gamma \vdash M^\circ : A$ and $\Gamma \vdash M^{\circ\circ} : A$.*

Proof. By induction on the structure of M , using corollary 5.4.

□

A term M is in *quasi-normal form* if only expansion rules at the *root position* are applicable to it and M is in *normal form* if no rule is applicable to it. So, every normal form is in quasi-normal form, while the converse does not necessarily hold.

Lemma 5.6

1. If M is in normal form, then $M^\circ = M$
2. If M is in quasi-normal form, then $M^{\circ\circ} = M$

Proof. By induction on the structure of M .

- $M \equiv *$.
 1. $*^\circ = *$.
 2. The property vacuously holds because $*$ is a normal form.
- $M \equiv x$.
 1. Since x is in normal form, it has neither a functional, nor a product, nor the \mathbf{T} type and then Δ_A is empty, where A is the type of x . Then $x^\circ = x$.
 2. $x^{\circ\circ} = x$ by definition.
- $M \equiv \lambda x : A.P$.
 1. Since M is in normal form, P is also in normal form and by induction hypothesis $P^\circ = P$. We have $(\lambda x : A.P)^\circ = \lambda x : A.P^\circ = \lambda x : A.P$.
 2. If $\lambda x : A.P$ is in quasi-normal form, it is also in normal form because we cannot apply an expansion rule to a *lambda-term*. By the previous paragraph $(\lambda x : A.P)^{\circ\circ} = \lambda x : A.P$.
- $M \equiv \langle P, Q \rangle$.
 1. Since M is in normal form, P and Q are also in normal form and by induction hypothesis $P^\circ = P$ and $Q^\circ = Q$. We have $\langle P, Q \rangle^\circ = \langle P^\circ, Q^\circ \rangle = \langle P, Q \rangle$.
 2. If $\langle P, Q \rangle$ is in quasi-normal form, it is also in normal form because we cannot apply an expansion rule to a pair. By the previous paragraph $\langle P, Q \rangle^{\circ\circ} = \langle P, Q \rangle$.
- $M \equiv (\text{rec } y : A.P)^i$.
 - If $i = 0$, then
 1. Since M is in normal form, P is also in normal form and by induction hypothesis $P^\circ = P$. On the other hand, M has neither a functional, nor a product, nor the \mathbf{T} type and then Δ_A is empty, where A is the type of M . We have $(\text{rec } y : A.P)^{0\circ} = \Delta_A^k(\text{rec } y : A.P^\circ)^0 = (\text{rec } y : A.P)^0$.
 2. Since M is in quasi normal form, P is in normal form and by induction hypothesis $P^\circ = P$. Then $(\text{rec } y : A.P)^{0\circ} = (\text{rec } y : A.P^\circ)^0 = (\text{rec } y : A.P)^0$.
 - If $i > 0$, then
 1. The property vacuously holds because $(\text{rec } y : A.P)^i$ is not in normal form.
 2. The property vacuously holds because $(\text{rec } y : A.P)^i$ is not in quasi-normal form.
- $M \equiv (PQ)$.

1. Suppose A is the type of M . Since M is in normal form, A is neither a functional, nor a product, nor the \mathbf{T} type and so Δ_A is empty. On the other hand P is in quasi-normal form and Q is in normal form, so by induction hypothesis $P^{\circ\circ} = P$ and $Q^\circ = Q$. We have $(PQ)^\circ = \Delta_A^k(P^{\circ\circ}Q^\circ) = (PQ)$.
 2. Since M is in quasi-normal form, P is in quasi-normal form and Q is in normal form and by induction hypothesis $P^{\circ\circ} = P$ and $Q^\circ = Q$. We have $(PQ)^{\circ\circ} = (P^{\circ\circ}Q^\circ) = (PQ)$.
- $M \equiv \text{Case}(P, R, N)$.
 1. Suppose A is the type of M . Since M is in normal form, A is neither a functional, nor a product, nor the \mathbf{T} type and so Δ_A is empty. On the other hand P, R and N are in normal form and by induction hypothesis $P^\circ = P$ and $R^\circ = R$ and $N^\circ = R$. We have $\text{Case}(P, R, N)^\circ = \Delta_A^k \text{Case}(P^\circ, R^\circ, N^\circ) = \text{Case}(P, R, N)$.
 2. Since M is in quasi-normal form, P, R and N are in normal form and by induction hypothesis $P^\circ = P$ and $R^\circ = R$ and $N^\circ = R$. We have $\text{Case}(P, R, N)^{\circ\circ} = \text{Case}(P^\circ, R^\circ, N^\circ) = \text{Case}(P, R, N)$.
 - $M \equiv \pi_i(P)$, for $i = 1, 2$.
 1. Suppose A is the type of M . Since M is in normal form, A is neither a functional, nor a product, nor the \mathbf{T} type and so Δ_A is empty. On the other hand P is in quasi-normal form and by induction hypothesis $P^{\circ\circ} = P$. We have $\pi_i(P)^\circ = \Delta_A^k \pi_i(P^{\circ\circ}) = \pi_i(P)$.
 2. Since M is in quasi-normal form, P is also in quasi-normal form and by induction hypothesis $P^{\circ\circ} = P$. We have $\pi_i(P)^{\circ\circ} = \pi_i(P^{\circ\circ}) = \pi_i(P)$.
 - $M \equiv \text{in}_C^i(P)$, for $i = 1, 2$.
 1. Since M is in normal form, P is also in normal form and by induction hypothesis $P^\circ = P$. We have $\text{in}_C^i(P)^\circ = \text{in}_C^i(P^\circ) = \text{in}_C^i(P)$.
 2. $\text{in}_C^i(P)$ in quasi-normal form implies $\text{in}_C^i(P)$ in normal form, and the property holds by the previous paragraph.

□

The next step is to prove that we can apply proposition 5.1 to our system, *i.e.*, for every one step reduction from M to N in the full system, there is a non empty reduction sequence in the system without expansions from any translation of M to a translation of N .

This lemma characterizes the reductions from a term $\Delta_{A \rightarrow B}^k M$ or $\Delta_{A \times B}^k M$ and is quite essential in all the properties shown in this section.

Lemma 5.7 *For any $k > 0$*

$$\Delta_{A \rightarrow B}^k M \Longrightarrow^+ \lambda w : A. \Delta_B^k (M(\Delta_A^k w))$$

and

$$\Delta_{A \times B}^k M \Longrightarrow^+ \langle \Delta_A^k \pi_1(M), \Delta_B^k \pi_2(M) \rangle$$

and the reduction sequences contain no expansion steps.

Proof. By induction on k .

If $k = 1$, then

$$\Delta_{A \rightarrow B} M \equiv (\lambda x : A \rightarrow B. \lambda w : A. \Delta_B(x(\Delta_A w))) M \xrightarrow{\beta} \lambda w : A. \Delta_B(M(\Delta_A w))$$

$$\Delta_{A \times B} M \equiv (\lambda x : A \times B. \langle \Delta_A \pi_1(x), \Delta_B \pi_2(x) \rangle) M \xrightarrow{\beta} \langle \Delta_A \pi_1(M), \Delta_B \pi_2(M) \rangle$$

Only the β -rule is used in this reduction.
 If $k > 1$, then

$$\begin{aligned}
& \Delta_{A \rightarrow B}^{k+1} M \\
& = \\
& \Delta_{A \rightarrow B}^k \Delta_{A \rightarrow B} M \\
& \Downarrow_{\beta} \\
& \Delta_{A \rightarrow B}^k \lambda w : A. \Delta_B (M(\Delta_A w)) \\
& \Downarrow_{+} \text{ by induction hypothesis (and without expansions steps)} \\
& \lambda w : A. \Delta_B^k (\lambda w : A. \Delta_B (M(\Delta_A w)) (\Delta_A^k w)) \\
& \Downarrow_{\beta} \\
& \lambda w : A. \Delta_B^k (\Delta_B (M(\Delta_A \Delta_A^k w))) \\
& = \\
& \lambda w : A. \Delta_B^{k+1} (M(\Delta_A^{k+1} w)) \\
& \\
& \Delta_{A \times B}^{k+1} M \\
& = \\
& \Delta_{A \times B}^k (\Delta_{A \times B} M) \\
& \Downarrow_{\beta} \\
& \Delta_{A \times B}^k \langle \Delta_A \pi_1 (M), \Delta_B \pi_2 (M) \rangle \\
& \Downarrow_{+} \text{ by induction hypothesis (and without expansion steps)} \\
& \langle \Delta_A^k \pi_1 (\langle \Delta_A \pi_1 (M), \Delta_B \pi_2 (M) \rangle), \Delta_B^k \pi_2 (\langle \Delta_A \pi_1 (M), \Delta_B \pi_2 (M) \rangle) \rangle \\
& \Downarrow_{\pi_1, \pi_2} \\
& \langle \Delta_A^k \Delta_A \pi_1 (M), \Delta_B^k \Delta_B \pi_2 (M) \rangle \\
& = \\
& \langle \Delta_A^{k+1} \pi_1 (M), \Delta_B^{k+1} \pi_2 (M) \rangle
\end{aligned}$$

□

We use N^{\otimes} to denote either N° or $N^{\circ\circ}$. In particular, $\overline{N^{\otimes}}$ will stand for a sequence of mixed N_i° 's and $N_i^{\circ\circ}$'s.

Lemma 5.8 *If $\Gamma \vdash M : A$, then*

1. $\exists k \geq 0, M^{\circ\circ}[\overline{z^{\otimes}}/\overline{x}] \Longrightarrow^* \Delta_A^k (M[\overline{z}/\overline{x}])^{\circ\circ}$
2. $\forall k \geq 0, \Delta_A^k M^{\circ}[\overline{z^{\otimes}}/\overline{x}] \Longrightarrow^* (M[\overline{z}/\overline{x}])^{\circ}$

and no expansions are performed in these reduction sequences.

Proof. We show the two properties by induction on the structure of M . More precisely, for the first statement we analyze each case, while for the second one it is enough to analyze those expressions M such that $M^{\circ} = M^{\circ\circ}$. Indeed, once we have already shown the first statement, the second can be easily shown in the following way for the expressions M such that $M^{\circ} = \Delta_A^h M^{\circ\circ}$ (for $h > 0$):

$$\begin{aligned}
& \Delta_A^k M^\circ[z^\otimes/\bar{x}] = \text{for some } h > 0 \\
& \Delta_A^k \Delta_A^h M^{\circ\circ}[z^\otimes/\bar{x}] \implies^* \text{ by the first statement} \\
& \Delta_A^{k+h} \Delta_A^m M[\bar{z}/\bar{x}]^{\circ\circ} = \\
& M[\bar{z}/\bar{x}]^\circ, \text{ because } h > 0
\end{aligned}$$

Every reduction built in the following proof contains no expansion steps, as it is constructed from one-step reductions that are no=3(e)-0.460706(60706(p))-0.1280Td[(a)0.433749(n)-0.575882(s)-0.774378(i)-0.338178(o)0.4

Z

$$\Delta_{B \rightarrow A}^h P[\bar{z}/\bar{x}]^{\circ\circ} Q^\circ[\bar{z}^\otimes/\bar{x}]$$

\Downarrow_+ by lemma 5.7

$$(\lambda w : B. \Delta_A^h (P[\bar{z}/\bar{x}]^{\circ\circ} (\Delta_B^h w))) Q^\circ[\bar{z}^\otimes/\bar{x}]$$

\Downarrow_β

$$\Delta_A^h (P[\bar{z}/\bar{x}]^{\circ\circ} (\Delta_B^h (Q^\circ[\bar{z}^\otimes/\bar{x}])))$$

\Downarrow_* by induction hypothesis

$$\begin{aligned} & \Delta_A^h (P[\bar{z}/\bar{x}]^{\circ\circ} Q[\bar{z}/\bar{x}]^\circ) \\ &= \\ & \Delta_A^h (P[\bar{z}/\bar{x}] Q[\bar{z}/\bar{x}]^{\circ\circ}) \\ &= \\ & \Delta_A^h (PQ[\bar{z}/\bar{x}]^{\circ\circ}) \end{aligned}$$

- $M \equiv \lambda y : B.P.$

1.

$$\begin{aligned} & (\lambda y : B.P)^{\circ\circ}[\bar{z}^\otimes/\bar{x}] \\ &= \\ & (\lambda y : B.P^\circ)[\bar{z}^\otimes/\bar{x}] \\ &= \\ & \lambda y : B.P^\circ[\bar{z}^\otimes/\bar{x}] \end{aligned}$$

\Downarrow_* by induction hypothesis

$$\begin{aligned} & \lambda y : B.P[\bar{z}/\bar{x}]^\circ \\ &= \\ & (\lambda y : B.P[\bar{z}/\bar{x}])^{\circ\circ} \\ &= \\ & \Delta_{B \rightarrow C}^0 ((\lambda y : B.P)[\bar{z}/\bar{x}])^{\circ\circ} \end{aligned}$$

2.

$$\begin{aligned}
& \Delta_{B \rightarrow C}^k (\lambda y : B.P)^\circ [\bar{z}^\otimes / \bar{x}] \\
& = \\
& \Delta_{B \rightarrow C}^k (\lambda y : B.P^\circ) [\bar{z}^\otimes / \bar{x}] \\
& = \\
& \Delta_{B \rightarrow C}^k \lambda y : B.P^\circ [\bar{z}^\otimes / \bar{x}]
\end{aligned}$$

\Downarrow_+ by lemma 5.7

$$\lambda w : B.\Delta_B^k ((\lambda y : B.P^\circ [\bar{z}^\otimes / \bar{x}]) (\Delta_C^k w))$$

\Downarrow_β

$$\lambda w : B.\Delta_B^k P^\circ [\bar{z}^\otimes / \bar{x}] [w^\circ / y]$$

\Downarrow_* by induction hypothesis

$$\begin{aligned}
& \lambda w : B.(P[\bar{z}/\bar{x}][w/y])^\circ \\
& = \\
& (\lambda w : B.P[\bar{z}/\bar{x}][w/y])^\circ \\
& =_\alpha \\
& (\lambda y : B.P[\bar{z}/\bar{x}])^\circ \\
& = \\
& ((\lambda y : B.P)[\bar{z}/\bar{x}])^\circ
\end{aligned}$$

- $M \equiv \pi_i(P)$, for $i = 1, 2$.

$$\begin{aligned}
& \pi_i(P)^\circ \circ [\bar{z}^\otimes / \bar{x}] \\
& = \\
& \pi_i(P^\circ \circ) [\bar{z}^\otimes / \bar{x}] \\
& = \\
& \pi_i(P^\circ \circ [\bar{z}^\otimes / \bar{x}])
\end{aligned}$$

\Downarrow_* by induction hypothesis

$$\pi_i(\Delta_{A_1 \times A_2}^h (P[\bar{z}/\bar{x}]^\circ \circ))$$

– If $h = 0$, then

$$\begin{aligned}
& \pi_i(P[\bar{z}/\bar{x}]^\circ \circ) \\
& = \\
& \pi_i(P[\bar{z}/\bar{x}])^\circ \circ \\
& = \\
& \Delta_{A_i}^0 \pi_i(P[\bar{z}/\bar{x}])^\circ \circ \\
& = \\
& \Delta_{A_i}^0 (\pi_i(P)[\bar{z}/\bar{x}])^\circ \circ
\end{aligned}$$

– If $h > 0$, then

$$\pi_i(\Delta_{A_1 \times A_2}^h(P[\bar{z}/\bar{x}])^{\circ\circ})$$

\Downarrow_+ by lemma 5.7

$$\pi_i(\langle \Delta_{A_1}^h \pi_1((P[\bar{z}/\bar{x}])^{\circ\circ}), \Delta_{A_2}^h \pi_2((P[\bar{z}/\bar{x}])^{\circ\circ}) \rangle)$$

\Downarrow_{π_i}

$$\Delta_{A_i}^h \pi_i((P[\bar{z}/\bar{x}])^{\circ\circ})$$

=

$$\Delta_{A_i}^h \pi_i(P[\bar{z}/\bar{x}]^{\circ\circ})$$

=

$$\Delta_{A_i}^h (\pi_i(P)[\bar{z}/\bar{x}])^{\circ\circ}$$

- $M \equiv in_C^i(P)$, for $i = 1, 2$ and then Δ_C is empty.

$$in_C^i(P)^{\circ\circ}[\bar{z}^{\otimes}/\bar{x}]$$

=

$$in_C^i(P^\circ)[\bar{z}^{\otimes}/\bar{x}]$$

=

$$in_C^i(P^\circ[\bar{z}^{\otimes}/\bar{x}])$$

\Downarrow_* by induction hypothesis

$$in_C^i(P[\bar{z}/\bar{x}]^\circ)$$

=

$$\Delta_A^0 in_C^i(P[\bar{z}/\bar{x}])^{\circ\circ}$$

=

$$\Delta_A^0 (in_C^i(P)[\bar{z}/\bar{x}])^{\circ\circ}$$

- $M \equiv \langle P, Q \rangle$.

1.

$$\langle P, Q \rangle^{\circ\circ}[\bar{z}^{\otimes}/\bar{x}]$$

=

$$\langle P^\circ, Q^\circ \rangle[\bar{z}^{\otimes}/\bar{x}]$$

=

$$\langle P^\circ[\bar{z}^{\otimes}/\bar{x}], Q^\circ[\bar{z}^{\otimes}/\bar{x}] \rangle$$

\Downarrow_* by induction hypothesis

$$\langle P[\bar{z}/\bar{x}]^\circ, Q[\bar{z}/\bar{x}]^\circ \rangle$$

=

$$\langle P[\bar{z}/\bar{x}], Q[\bar{z}/\bar{x}] \rangle^{\circ\circ}$$

=

$$\Delta_{A \times B}^0 \langle P[\bar{z}/\bar{x}], Q[\bar{z}/\bar{x}] \rangle^{\circ\circ}$$

=

$$\Delta_{A \times B}^0 (\langle P, Q \rangle[\bar{z}/\bar{x}])^{\circ\circ}$$

2.

$$\begin{aligned}
& \Delta_{A \times B}^k \langle P, Q \rangle^\circ [\bar{z}^\otimes / \bar{x}] \\
& = \\
& \Delta_{A \times B}^k \langle P^\circ, Q^\circ \rangle [\bar{z}^\otimes / \bar{x}] \\
& = \\
& \Delta_{A \times B}^k \langle P^\circ [\bar{z}^\otimes / \bar{x}], Q^\circ [\bar{z}^\otimes / \bar{x}] \rangle
\end{aligned}$$

– If $k = 0$, then

$$\begin{aligned}
& \langle P^\circ [\bar{z}^\otimes / \bar{x}], Q^\circ [\bar{z}^\otimes / \bar{x}] \rangle \\
& \Downarrow_* \text{ by induction hypothesis} \\
& \langle P[\bar{z}/\bar{x}]^\circ, Q[\bar{z}/\bar{x}]^\circ \rangle \\
& = \\
& \langle P[\bar{z}/\bar{x}], Q[\bar{z}/\bar{x}] \rangle^\circ \\
& = \\
& \langle \langle P, Q \rangle [\bar{z}/\bar{x}] \rangle^\circ
\end{aligned}$$

– If $k > 0$, then

$$\begin{aligned}
& \Delta_{A \times B}^k \langle P^\circ [\bar{z}^\otimes / \bar{x}], Q^\circ [\bar{z}^\otimes / \bar{x}] \rangle \\
& \Downarrow_+ \text{ by lemma 5.7} \\
& \langle \Delta_A^k \pi_1(P^\circ [\bar{z}^\otimes / \bar{x}] Q^\circ [\bar{z}^\otimes / \bar{x}]), \Delta_B^k \pi_2(\langle P^\circ [\bar{z}^\otimes / \bar{x}], Q^\circ [\bar{z}^\otimes / \bar{x}] \rangle) \rangle \\
& \Downarrow \pi_1, \pi_2 \\
& \langle \Delta_A^k P^\circ [\bar{z}^\otimes / \bar{x}], \Delta_B^k Q^\circ [\bar{z}^\otimes / \bar{x}] \rangle \\
& \Downarrow_* \text{ by induction hypothesis} \\
& \langle P[\bar{z}/\bar{x}]^\circ, Q[\bar{z}/\bar{x}]^\circ \rangle \\
& = \\
& \langle P[\bar{z}/\bar{x}], Q[\bar{z}/\bar{x}] \rangle^\circ \\
& = \\
& \langle \langle P, Q \rangle [\bar{z}/\bar{x}] \rangle^\circ
\end{aligned}$$

• $M \equiv (\text{rec } y : A.P)^i$.

$$\begin{aligned}
& (\text{rec } y : A.P)^{i^\circ\circ} [\bar{z}^\otimes / \bar{x}] \\
& = \\
& (\text{rec } y : A.P^\circ)^i [\bar{z}^\otimes / \bar{x}] \\
& \Downarrow_* \text{ by induction hypothesis} \\
& (\text{rec } y : A.(P[\bar{z}/\bar{x}])^\circ)^i \\
& = \\
& \Delta_A^0 (\text{rec } y : A.P[\bar{z}/\bar{x}])^{i^\circ\circ} \\
& = \\
& \Delta_A^0 ((\text{rec } y : A.P)^i [\bar{z}/\bar{x}])^\circ\circ
\end{aligned}$$

• $M \equiv \text{Case}(P, Q, R)$.

$$\begin{aligned}
& \text{Case}(P, Q, R)^{\circ\circ}[\overline{z^\otimes}/\overline{x}] \\
& = \\
& \text{Case}(P^\circ, Q^\circ, R^\circ)[\overline{z^\otimes}/\overline{x}] \\
& = \\
& \text{Case}(P^\circ[\overline{z^\otimes}/\overline{x}], Q^\circ[\overline{z^\otimes}/\overline{x}], R^\circ[\overline{z^\otimes}/\overline{x}]) \\
& \Downarrow_* \text{ by induction hypothesis} \\
& \text{Case}(P[\overline{z}/\overline{x}]^\circ, Q[\overline{z}/\overline{x}]^\circ, R[\overline{z}/\overline{x}]^\circ) \\
& = \\
& \text{Case}(P[\overline{z}/\overline{x}], Q[\overline{z}/\overline{x}], R[\overline{z}/\overline{x}])^{\circ\circ} \\
& = \\
& (\text{Case}(P, Q, R)[\overline{z}/\overline{x}])^{\circ\circ} \\
& = \\
& \Delta_A^0(\text{Case}(P, Q, R)[\overline{z}/\overline{x}])^{\circ\circ}
\end{aligned}$$

□

Corollary 5.9 *If $\Gamma \vdash M : A$, then*

$$\forall k \geq 0, \Delta_A^k M^\circ \Longrightarrow^* M^\circ$$

and no expansions are performed in the reduction sequences.

The following property is essential to show that every time we perform a β -reduction on a term M in the original system, any translation of M reduces to a translation of the term we have obtained via \rightarrow_β from M . Take for example the reduction $(\lambda x : A.M)N \rightarrow_\beta M[N/x]$. We know that $((\lambda x : A.M)N)^\circ = \Delta_A^k((\lambda x : A.M^\circ)N^\circ)$ and we want to show that there is a *non empty* reduction sequence leading to $M[N/x]^\circ$. Since $\Delta_A^k((\lambda x : A.M^\circ)N^\circ) \rightarrow_\beta \Delta_A^k M^\circ[N^\circ/x]$, we have now to check that the term $(M[N/x])^\circ$ can be reached. We state the property as follows:

Lemma 5.10 *If $\Gamma \vdash M : A$, then*

1. $\forall k \geq 0, \Delta_A^k M^\circ[\overline{N^\circ}/\overline{x}] \Longrightarrow^* (M[\overline{N}/\overline{x}])^\circ$
2. $\exists k \geq 0, M^{\circ\circ}[\overline{N^\circ}/\overline{x}] \Longrightarrow^* \Delta_A^k (M[\overline{N}/\overline{x}])^{\circ\circ}$

and no expansions are performed in the reduction sequences.

Proof. The proof of this lemma can be done exactly as for lemma 5.8, except that now we have to use corollary 5.9 in two points: to show that

$$\Delta_A^{k+h} \Delta_A^m M[\overline{N}/\overline{x}]^{\circ\circ} \Longrightarrow^* M[\overline{N}/\overline{x}]^\circ$$

if $M[\overline{N}/\overline{x}]$ is a pair or a λ -abstraction, and in the case $M \equiv x_i$, where to prove the first statement we need to proceed as follows:

$$\Delta_A^k x_i^\circ[\overline{N^\circ}/\overline{x}] = \Delta_A^k (\Delta_A^m x_i)[\overline{N^\circ}/\overline{x}] = \Delta_A^{k+m} N_i^\circ.$$

$$\text{By corollary 5.9, } \Delta_A^{k+m} N_i^\circ \Longrightarrow^* N_i^\circ = (x_i[\overline{N}/\overline{x}])^\circ.$$

Notice that there are no expansions in the sequence $\Delta_A^{k+m} N_i^\circ \Longrightarrow^* N_i^\circ$ by corollary 5.9.

□

Lemma 5.11 *If $\Gamma \vdash M : A$, then*

1. $\forall k \geq 0, \Delta_A^k M^\circ[\overline{N^{\circ\circ}}/\overline{x}] \Longrightarrow^* (M[\overline{N}/\overline{x}])^\circ$
2. $\exists k \geq 0, M^{\circ\circ}[\overline{N^{\circ\circ}}/\overline{x}] \Longrightarrow^* \Delta_A^k (M[\overline{N}/\overline{x}])^{\circ\circ}$

and no expansions are performed in the reduction sequences.

Proof. The proof of this lemma can be done exactly as for lemma 5.8, except that now we have to use corollary 5.9 in two points: to show that

$$\Delta_A^{k+h} \Delta_A^m M[\overline{N}/\overline{x}]^{\circ\circ} \Longrightarrow^* M[\overline{N}/\overline{x}]^\circ$$

if $M[\overline{N}/\overline{x}]$ is a pair or a λ -abstraction, and in the case $M \equiv x_i$, where to prove the first statement we need to proceed as follows:

$$\Delta_A^k x_i^\circ[\overline{N}^{\circ\circ}/\overline{x}] = \Delta_A^k (\Delta_A^m x_i) [\overline{N}^{\circ\circ}/\overline{x}] = \Delta_A^{k+m} N_i^{\circ\circ}.$$

If $N_i^\circ = N_i^{\circ\circ}$, then by corollary 5.9 $\Delta_A^{k+m} N_i^{\circ\circ} \Longrightarrow^* N_i^\circ = (x_i[\overline{N}/\overline{x}])^\circ$.

In the other case $\Delta_A^{k+m} N_i^{\circ\circ} = N_i^\circ = (x_i[\overline{N}/\overline{x}])^\circ$.

Notice that there are no expansions in the sequence $\Delta_A^{k+m} N_i^{\circ\circ} \Longrightarrow^* N_i^\circ$ by corollary 5.9.

□

Lemma 5.12 *If $M \xrightarrow{\eta, \delta, Top} N$, then $M \xrightarrow{\sigma, \delta, \neg\eta} N^\circ$.*

Proof.

- If M is of type $A \times B$ and $M \xrightarrow{\delta} \langle \pi_1(M), \pi_2(M) \rangle$

We know that $\exists k > 0$ such that $M^\circ = \Delta_{A \times B}^k M^{\circ\circ}$. By corollary 5.7

$$\Delta_{A \times B}^k M^{\circ\circ} \Longrightarrow^+ \langle \Delta_A^k \pi_1(M^{\circ\circ}), \Delta_B^k \pi_2(M^{\circ\circ}) \rangle$$

and the sequence has no expansion rules.

The last term is equal to $\langle \pi_1(M)^\circ, \pi_2(M)^\circ \rangle = \langle \pi_1(M), \pi_2(M) \rangle^\circ$ and then the property holds.

- If M is of type $A \rightarrow B$ and $M \xrightarrow{\eta} \lambda y : A.M y$

We know that $\exists k > 0$ such that $M^\circ = \Delta_{A \rightarrow B}^k M^{\circ\circ}$. By corollary 5.7 $\Delta_{A \rightarrow B}^k M^{\circ\circ} \Longrightarrow^+ \lambda y : A. \Delta_B^k (M^{\circ\circ}(\Delta_A^k y))$ and the the sequence has no expansion rules.

The last term is equal to $\lambda y : A. \Delta_B^k (M^{\circ\circ} y^\circ) = \lambda y : A. (M y)^\circ = (\lambda y : A. M y)^\circ$ and then the property holds.

- If $M : \mathbf{T}$ and $M \xrightarrow{Top} *$.

By lemma 5.5 $M^\circ : \mathbf{T}$ and so $M^\circ \xrightarrow{Top} * = *^\circ$.

□

Using 5.10 we can show now:

Theorem 5.13 (Simulation) *If $\Gamma \vdash M : A$ and $M \Longrightarrow N$, then*

1. $\exists k \geq 0$ such that $M^{\circ\circ} \Longrightarrow^+ \Delta_A^k N^{\circ\circ}$ if $M \xrightarrow{\neg\eta, \neg\delta} N$

2. $M^\circ \Longrightarrow^+ N^\circ$

and there are no expansions in these reduction sequences.

Proof. We show the property by induction on the structure of M . More precisely, for the first statement we analyze each case, while for the second there are two cases:

- if $M \xrightarrow{\eta, \delta} N$, then apply lemma 5.12
- if $M \xrightarrow{\neg\eta, \neg\delta} N$, then it is enough to analyze only the cases such that $M^\circ = M^{\circ\circ}$, because when $M^\circ = \Delta_A^h M^{\circ\circ}$ (for $h > 0$) we have easily:

$$M^\circ = \Delta_A^h M^{\circ\circ} \Longrightarrow^+ \text{(by the first statement)} \Delta_A^h \Delta_A^k N^{\circ\circ} = \Delta_A^{h+k} N^{\circ\circ}$$

Then either N is not a pair or a λ -abstraction, which gives $\Delta_A^{h+k} N^{\circ\circ} = N^\circ$ because $h > 0$, or otherwise $\Delta_A^{h+k} N^{\circ\circ} = \Delta_A^{h+k} N^\circ \Longrightarrow^* N^\circ$ by lemma 5.9.

In order to conclude that the reductions in the statements of the lemma contain no expansions, it suffices to notice that every reduction built in the following proof contains no expansion steps: indeed it is constructed from one-step reductions that are not expansions or from reductions obtained by induction hypothesis (and thus without expansions) or from reductions obtained by lemma 5.12, lemma 5.7, (again without expansions).

Now, we can analyze the cases involved in the proof of the first and the second statement.

- $M \equiv *$. It is in normal form.
- $M \equiv x$. The only possible case is $x \xrightarrow{Top} *$, where $x : \mathbf{T}$. Then, $x^{\circ\circ} = x \xrightarrow{Top} * = \Delta_{\mathbf{T}}^0 *^{\circ\circ}$.
- $M \equiv (P_1 Q_1)$.

– If $(P_1 Q_1) : \mathbf{T}$ and $(P_1 Q_1) \xrightarrow{Top} *$, then $(P_1 Q_1)^{\circ\circ} : \mathbf{T}$ by lemma 5.5 and then

$$(P_1 Q_1)^{\circ\circ} \xrightarrow{Top} * = \Delta_{\mathbf{T}}^0 *^{\circ\circ}$$

– If $(\lambda x : C.R)Q_1 \xrightarrow{\beta} R[Q_1/x]$.

$$\begin{aligned} ((\lambda x : C.R)Q_1)^{\circ\circ} &= \\ ((\lambda x : C.R)^{\circ\circ} Q_1^{\circ}) &= \\ ((\lambda x : C.R^{\circ})Q_1^{\circ}) &\xrightarrow{\beta} \\ R^{\circ}[Q_1^{\circ}/x] &\xRightarrow{*} \text{ (by lemma 5.10)} \\ R[Q_1/x]^{\circ} &= \\ \Delta_A^h R[Q_1/x]^{\circ\circ} & \end{aligned}$$

– If $(P_1 Q_1) \Longrightarrow (P_2 Q_1)$, where $P_1 \Longrightarrow P_2$.

Since $P_1 \xrightarrow{\neg\eta} P_2$ because $(P_1 Q_1) \Longrightarrow (P_2 Q_1)$, we have by induction hypothesis a reduction sequence $P_1^{\circ\circ} \xRightarrow{+} \Delta_{B \rightarrow A}^h P_2^{\circ\circ}$ without expansions. Then

$$\begin{aligned} (P_1 Q_1)^{\circ\circ} &= \\ P_1^{\circ\circ} Q_1^{\circ} &\xRightarrow{+} \text{ by induction hypothesis since } P_1 \xrightarrow{\neg\eta} P_2 \\ (\Delta_{B \rightarrow A}^h P_2^{\circ\circ}) Q_1^{\circ} & \end{aligned}$$

If $h = 0$, then $(P_2^{\circ\circ} Q_1^{\circ}) = (P_2 Q_1)^{\circ\circ}$.

If $h > 0$, then

$$\begin{aligned} &(\Delta_{B \rightarrow A}^h P_2^{\circ\circ}) Q_1^{\circ} \\ &\Downarrow_+ \text{ by lemma 5.7} \\ &(\lambda w : B. \Delta_A^h (P_2^{\circ\circ} (\Delta_B^h w))) Q_1^{\circ} \\ &\downarrow \beta \\ &\Delta_A^h (P_2^{\circ\circ} (\Delta_B^h Q_1^{\circ})) \\ &\Downarrow * \text{ by corollary 5.9} \\ &\Delta_A^h (P_2^{\circ\circ} Q_1^{\circ}) \\ &= \\ &\Delta_A^h (P_2 Q_1)^{\circ\circ} \end{aligned}$$

– If $\langle P_1 Q_1 \rangle \Longrightarrow \langle P_1 Q_2 \rangle$, where $Q_1 \Longrightarrow Q_2$

$$\begin{aligned} & \langle P_1 Q_1 \rangle^{\circ\circ} = \\ & P_1^{\circ\circ} Q_1^{\circ} \Longrightarrow^+ \text{ by induction hypothesis} \\ & P_1^{\circ\circ} Q_2^{\circ} = \\ & \langle P_1 Q_2 \rangle^{\circ\circ} \end{aligned}$$

• $M \equiv \langle P_1, Q_1 \rangle$

– If $\langle P_1, Q_1 \rangle \Longrightarrow \langle P_2, Q_1 \rangle$, where $P_1 \Longrightarrow P_2$.

1.

$$\begin{aligned} & \langle P_1, Q_1 \rangle^{\circ\circ} \\ & \langle P_1^{\circ}, Q_1^{\circ} \rangle \Longrightarrow^+ \text{ by induction hypothesis} \\ & \langle P_2^{\circ}, Q_1^{\circ} \rangle = \\ & \langle P_2, Q_1 \rangle^{\circ\circ} = \\ & \Delta_A^0 \langle P_2, Q_1 \rangle^{\circ\circ} \end{aligned}$$

2. Since $\langle P, Q \rangle^{\circ\circ} = \langle P, Q \rangle^{\circ}$, we have $\langle P_1, Q_1 \rangle^{\circ} \Longrightarrow^+ \langle P_2, Q_1 \rangle^{\circ}$ by the previous statement.

– If $\langle P_1, Q_1 \rangle \Longrightarrow \langle P_1, Q_2 \rangle$, where $Q_1 \Longrightarrow Q_2$.

1.

$$\begin{aligned} & \langle P_1, Q_1 \rangle^{\circ\circ} \\ & \langle P_1^{\circ}, Q_1^{\circ} \rangle \Longrightarrow^+ \text{ by induction hypothesis} \\ & \langle P_1^{\circ}, Q_2^{\circ} \rangle = \\ & \langle P_1, Q_2 \rangle^{\circ\circ} = \\ & \Delta_A^0 \langle P_1, Q_2 \rangle^{\circ\circ} \end{aligned}$$

2. Since $\langle P, Q \rangle^{\circ\circ} = \langle P, Q \rangle^{\circ}$, we have $\langle P_1, Q_1 \rangle^{\circ} \Longrightarrow^+ \langle P_1, Q_2 \rangle^{\circ}$ by the previous statement.

• $M \equiv \lambda x : A. P_1$

Then $\lambda x : A. P_1 \Longrightarrow \lambda x : A. P_2$, where $P_1 \Longrightarrow P_2$.

1.

$$\begin{aligned} & (\lambda x : A. P_1)^{\circ\circ} \\ & \lambda x : A. P_1^{\circ} \Longrightarrow^+ \text{ by induction hypothesis} \\ & \lambda x : A. P_2^{\circ} = \\ & \Delta_{A \rightarrow B}^0 (\lambda x : A. P_2)^{\circ\circ} \end{aligned}$$

2. Since $(\lambda x : A. P_i)^{\circ} = (\lambda x : A. P_i)^{\circ\circ}$ we have $(\lambda x : A. P_1)^{\circ} \Longrightarrow^+ (\lambda x : A. P_2)^{\circ}$ by the previous statement.

• $M \equiv in_C^i(P_1)$, for $i = 1, 2$ where $P_1 \Longrightarrow P_2$.

Then $in_C^i(P_1) \Longrightarrow in_C^i(P_2)$

$$\begin{aligned} & in_C^i(P_1)^{\circ\circ} = \\ & in_C^i(P_1^{\circ}) \Longrightarrow^+ \text{ by induction hypothesis} \\ & in_C^i(P_2^{\circ}) = \\ & \Delta_{B+C}^0 in_C^i(P_2)^{\circ\circ} \end{aligned}$$

• $M \equiv \pi_i(P_1)$, for $i = 1, 2$.

– If $\pi_i(P_1) : \mathbf{T}$ and $\pi_i(P_1) \xrightarrow{Top} *$, then $\pi_i(P_1)^{\circ\circ} : \mathbf{T}$ by lemma 5.5 and $\pi_i(P_1)^{\circ\circ} \xrightarrow{Top} * = \Delta_{Top}^0 *^{\circ\circ}$.

- If $\pi_i(P_1) \Longrightarrow \pi_i(P_2)$, where $P_1 \Longrightarrow P_2$.

Since $P_1 \xrightarrow{-\delta} P_2$ because $\pi_i(P_1) \Longrightarrow \pi_i(P_2)$, we have by induction hypothesis a reduction sequence $P_1^{\circ\circ} \Longrightarrow^+ \Delta_{B \times A}^h P_2^{\circ\circ}$ without expansions. Then

$$\begin{aligned} \pi_i(P_1)^{\circ\circ} &= \\ \pi_i(P_1^{\circ\circ}) &\Longrightarrow^+ \text{ by induction hypothesis} \\ \pi_i(\Delta_{A_1 \times A_2}^h P_2^{\circ\circ}) & \end{aligned}$$

If $h = 0$, then $\pi_i(P_2^{\circ\circ}) = \pi_i(P_2)^\circ$.

If $h > 0$, then

$$\begin{aligned} &\pi_i(\Delta_{A_1 \times A_2}^h P_2^{\circ\circ}) \\ &\Downarrow_+ \text{ by lemma 5.7} \\ &\pi_i(\langle \Delta_{A_1}^h \pi_1(P_2^{\circ\circ}), \Delta_{A_2}^h \pi_2(P_2^{\circ\circ}) \rangle) \\ &\Downarrow_{\pi_i} \\ &\Delta_{A_i}^h \pi_i(P_2^{\circ\circ}) \\ &= \\ &\Delta_{A_i}^h \pi_i(P_2)^{\circ\circ} \end{aligned}$$

- $M \equiv \text{Case}(P_1, R_1, N_1)$

- If $\text{Case}(P_1, R_1, N_1) : \mathbf{T}$ and $\text{Case}(P_1, R_1, N_1) \xrightarrow{\text{Top}} *$, then $\text{Case}(P_1, R_1, N_1)^{\circ\circ} : \mathbf{T}$ by lemma 5.5 and $\pi_i(P_1)^{\circ\circ} \xrightarrow{\text{Top}} * = \Delta_{\text{Top}}^0 *^{\circ\circ}$.

- $\text{Case}(in_{C_1+C_2}^i(S), R_1, R_2) \xrightarrow{\rho} R_i S$

$$\begin{aligned} &\text{Case}(in_{C_1+C_2}^i(S), R_1, R_2)^{\circ\circ} = \\ &\text{Case}(in_{C_1+C_2}^i(S)^\circ, R_1^\circ, R_2^\circ) = \\ &\text{Case}(in_{C_1+C_2}^i(S)^\circ, R_1^\circ, R_2^\circ) \xrightarrow{\rho} \\ &R_1^\circ S^\circ = \\ &(\Delta_{C_i \rightarrow A}^h R_1^{\circ\circ}) S^\circ \Longrightarrow^+ \text{ by lemma ??} \\ &(\lambda w : C_i. \Delta_A^h (R_1^{\circ\circ} (\Delta_{C_i}^h w))) S^\circ \xrightarrow{\beta} \\ &\Delta_A^h (R_1^{\circ\circ} (\Delta_{C_i}^h S^\circ)) \Longrightarrow^* \text{ by corollary 5.9} \\ &\Delta_A^h (R_1^{\circ\circ} S^\circ) = \\ &\Delta_A^h (R_1 S)^{\circ\circ} \end{aligned}$$

- $\text{Case}(P_1, R_1, N_1) \Longrightarrow \text{Case}(P_2, R_1, N_1)$, where $P_1 \Longrightarrow P_2$.

$$\begin{aligned} &\text{Case}(P_1, R_1, N_1)^{\circ\circ} = \\ &\text{Case}(P_1^\circ, R_1^\circ, N_1^\circ) \Longrightarrow^+ \text{ by induction hypothesis since} \\ &\text{Case}(P_2^\circ, R_1^\circ, N_1^\circ) = \\ &\text{Case}(P_2, R_1, N_1)^{\circ\circ} = \\ &\Delta_A^0 \text{Case}(P_2, R_1, N_1)^{\circ\circ} \end{aligned}$$

- $\text{Case}(P_1, R_1, N_1) \Longrightarrow \text{Case}(P_1, R_2, N_1)$, where $R_1 \Longrightarrow R_2$

$$\begin{aligned} &\text{Case}(P_1, R_1, N_1)^{\circ\circ} = \\ &\text{Case}(P_1^\circ, R_1^\circ, N_1^\circ) \Longrightarrow^+ \text{ by induction hypothesis} \\ &\text{Case}(P_1^\circ, R_2^\circ, N_1^\circ) = \\ &\text{Case}(P_1, R_2, N_1)^{\circ\circ} = \\ &\Delta_A^0 \text{Case}(P_1, R_2, N_1)^{\circ\circ} \end{aligned}$$

– $Case(P_1, R_1, N_1) \Longrightarrow Case(P_1, R_1, N_2)$, where $N_1 \Longrightarrow N_2$

$$\begin{aligned} & Case(P_1, R_1, N_1)^{\circ\circ} = \\ & Case(P_1^\circ, R_1^\circ, N_1^\circ) \Longrightarrow^+ \text{ by induction hypothesis} \\ & Case(P_1^\circ, R_1^\circ, N_2^\circ) = \\ & Case(P_1, R_1, N_2)^{\circ\circ} = \\ & \Delta_A^0 Case(P_1, R_1, N_2)^{\circ\circ} \end{aligned}$$

• $M \equiv (rec\ y : B.P_1)^i$

– If $(rec\ y : \mathbf{T}.P_1)^i : \mathbf{T}$ and $(rec\ y : \mathbf{T}.P_1)^i \xrightarrow{Top} *$, then $((rec\ y : \mathbf{T}.P_1)^i)^{\circ\circ} : \mathbf{T}$ by lemma 5.5 and $(rec\ y : \mathbf{T}.P_1)^{i\circ\circ} \xrightarrow{Top} * = \Delta_{Top}^0 *^{\circ\circ}$.

– $(rec\ y : A.P_1)^i \xrightarrow{rec} P_1[(rec\ y : A.P_1)^{i-1}/y]$

$$\begin{aligned} & ((rec\ y : A.P_1)^i)^{\circ\circ} = \\ & (rec\ y : A.P_1^\circ)^i \xrightarrow{rec} \\ & P_1^\circ[(rec\ y : A.P_1^\circ)^{i-1}/y] = \\ & P_1^\circ[(rec\ y : A.P_1)^{i-1\circ\circ}/y] \Longrightarrow^* \text{ by lemma 5.8} \\ & (P_1[(rec\ y : A.P_1)^{i-1}/y])^\circ = \\ & \Delta_A^h (P_1[(rec\ y : A.P_1)^{i-1}/y])^{\circ\circ} \end{aligned}$$

– $(rec\ y : A.P_1)^i \Longrightarrow (rec\ y : A.P_2)^i$.

$$\begin{aligned} & ((rec\ y : A.P_1)^i)^{\circ\circ} = \\ & (rec\ y : A.P_1^\circ)^i \Longrightarrow^+ \text{ by induction hypothesis} \\ & (rec\ y : A.P_2^\circ)^i = \\ & \Delta_A^0 (rec\ y : A.P_2)^{i\circ\circ} \end{aligned}$$

□

5.2 Strong Normalization of the Full Calculus

Having shown that our translation satisfies the hypothesis of Proposition 5.1, all we are now left to prove is that the bounded reduction system without expansion rules is strongly normalizing. This can be established by one of the standard techniques of reducibility, and does not present essential difficulties once the right definitions of *stability* or *reducibility* are given. In 6 and 7 we provide two proofs, one adapting the proof provided by Poigné and Voss in [PV87], and the other adapting Girard's proof from [GLT90]. It is then finally possible to state the following

Theorem 5.14 (Strong normalization)

The reduction \Longrightarrow for the bounded system with expansions is strongly normalizing.

Proof. By proposition 5.1, theorem 5.13 and Corollary 6.11 (or 7.7). □

6 Strong Normalization via stability

We have shown that every reduction sequence starting at M involving the rules presented in section 3 can be simulated by the translation of M in the same calculus but without expansions. We will show in this section that the calculus without expansions is strongly normalizing.

6.1 Stability

We define a set of *stable terms of type A* by induction on the type A in the following way:

- For M of atomic type A , M is stable if and only if it is strongly normalizing.
- For M of type $A_1 \times A_2$, M is stable if and only if it is strongly normalizing and whenever M reduces to $\langle M_1, M_2 \rangle$, then M_1 and M_2 are stable terms of type A_1 and A_2 respectively.
- For M of type $A_1 + A_2$, M is stable if and only if it is strongly normalizing and whenever M reduces to $in^i_{A_1+A_2}(M')$ then M' is stable of type A_i .
- For M of type $A_1 \rightarrow A_2$, M is stable if and only if for every stable term N of type A_1 , MN is a stable term of type A_2 .

Proposition 6.1 *Let M be a term of type $A \rightarrow B$*

1. *if M is stable, then $MQ_1 \dots Q_k$ is stable for arbitrary stable $Q_1 \dots Q_k$ of appropriate types,*
2. *if $MQ_1 \dots Q_k$ is stable for arbitrary stable $Q_1 \dots Q_k$ of appropriate types, then M is stable.*

Proof.

By induction on k . For $k = 1$, this is just our definition of stability, while for $k > 1$:

1. we know by induction hypothesis that $MQ_1 \dots Q_{k-1}$ is stable, and then $MQ_1 \dots Q_k$ is stable by definition since Q_k is stable too.
2. if $MQ_1 \dots Q_k$ is stable for arbitrary stable terms $Q_1 \dots Q_k$, then by definition of stability, $MQ_1 \dots Q_{k-1}$ is stable for arbitrary $Q_1 \dots Q_{k-1}$, and then by induction hypothesis M is stable.

□

Remark 6.2 *We can then use equivalently as a definition for stability $k > 1$ or $k = 1$. We will use in the following the most suitable one for each case.*

Notation 6.3 *In what follows we will use often \vec{Q} to denote a sequence of terms $Q_1 \dots Q_k$, with the convention that $M\vec{Q}$ really stands for $MQ_1 \dots Q_k$.*

Lemma 6.4

- *If M, N, \vec{Q} are strongly normalizing, then the terms $x\vec{Q}, Mx, in^i_A(M), (rec\ y : B.M)^0\vec{Q}$ and $\langle M, N \rangle$ are strongly normalizing.*
- *If M, \vec{Q} are strongly normalizing and M has product type but does not reduce to a pair, the $\pi_i(M)\vec{Q}$ is strongly normalizing.*
- *If P, M, N, \vec{Q} are strongly normalizing and P has sum type but does not reduce to in^i , then $Case(P, M, N)\vec{Q}$ is strongly normalizing.*

Proof. The argument of the proof is the same for all the cases: we show that a generic reduction sequence starting from the given term always terminates. We assume in what follows that no *Top* rule is applied at the root of our term during the reduction, as in this case the reduction sequence simply stops and is then clearly finite.

- $- x\vec{Q}$. There are no expansions in this system, so a reduction starting at $x\vec{Q}$ can only proceed in the Q'_i s. Since the \vec{Q} are strongly normalizing, this means that the reduction sequence also terminates, so we conclude that $x\vec{Q}$ is strongly normalizing.

- Mx . Consider a given reduction sequence starting from Mx . There are two possibilities:
 - * M does not reduce to a λ -expression in the reduction sequence. Then, the reduction sequence starting at Mx terminates because it is of the form $Mx \Rightarrow M_1x \Rightarrow \dots \Rightarrow M_nx$, where $M \Rightarrow M_1 \Rightarrow \dots \Rightarrow M_n$ is a terminating reduction sequence starting at M .
 - * M reduces at some point to $\lambda y : B.P$. Then the reduction sequence starting at Mx is of the form $Mx \Rightarrow^* (\lambda y : B.P)x' \Rightarrow^* (\lambda y : B.P')x'' \Rightarrow^* P'[x''/y] \Rightarrow^* \dots$, where $P \Rightarrow^* P'$ and $x \Rightarrow^* x' \Rightarrow^* x''$ (because x can reduce to $*$). On the other hand P' is strongly normalizing as P is, and thus also $P'[x/y]$ and $P'[* /y]$ must be strongly normalizing⁶. Therefore the reduction sequence is terminating.

We conclude that Mx is strongly normalizing.

- $in_{A_1+A_2}^i(M)$. A reduction sequence starting at $in_{A_1+A_2}^i(M)$ is of the form $in_{A_1+A_2}^i(M) \Rightarrow in_{A_1+A_2}^i(M_1) \Rightarrow in_{A_1+A_2}^i(M_2) \Rightarrow \dots$, where $M \Rightarrow M_1 \Rightarrow M_2 \Rightarrow \dots$ since the leftmost in^i cannot be removed. Since M is strongly normalizing, these two sequences are both finite.
- $(rec\ y : B.M)^0 \vec{Q}$. Since the term $(rec\ y : B.M)^0$ cannot be reduced at the root position (neither by a *rec*-rule nor by a *Top*-rule), then only internal reductions in M are possible. Therefore a reduction starting at $(rec\ y : B.M)^0 \vec{Q}$ only proceeds in M, \vec{Q} that are strongly normalizing by hypothesis. Hence, the reduction sequence is finite.
- $\langle M, N \rangle$. A reduction sequence starting at $\langle M, N \rangle$ is of the form $\langle M, N \rangle \Rightarrow^* \langle M_1, N_1 \rangle \Rightarrow^* \langle M_2, N_2 \rangle \Rightarrow^* \dots$, where $M \Rightarrow^* M_1 \Rightarrow^* M_2 \Rightarrow^* \dots$ and $N \Rightarrow^* N_1 \Rightarrow^* N_2 \Rightarrow^* \dots$. Since M and N are strongly normalizing, these reduction sequences are all finite.
- $\pi_i(M) \vec{Q}$. Since by hypothesis M does not reduce to a pair, the leftmost π_i cannot be removed and therefore a reduction starting at $\pi_i(M) \vec{Q}$ only proceeds in M, \vec{Q} that are strongly normalizing by hypothesis. Therefore the reduction sequence is finite.
- $Case(P, M, N) \vec{Q}$. Since by hypothesis P does not reduce to an in^i , the leftmost *Case* cannot be removed and therefore a reduction starting at $Case(P, M, N) \vec{Q}$ only proceeds in P, M, N, \vec{Q} that are strongly normalizing by hypothesis. Hence, the reduction sequence terminates.

□

6.2 Properties of stability

Lemma 6.5

1. Every stable term M of type A is strongly normalizing.
2. A term $x \vec{Q}$ of type A is stable for arbitrary strongly normalizing terms \vec{Q} .

Proof. By induction on the type A

- If A is not a functional type:
 1. By definition.
 2. By lemma 6.4 $x \vec{Q}$ is strongly normalizing. As pointed out in the proof of lemma 6.4, either $x \vec{Q}$ reduces to $*$, or reductions from $x \vec{Q}$ can only proceed in the Q_i 's. Therefore, $x \vec{Q}$ cannot reduce to a pair, nor to an in^i and thus $x \vec{Q}$ is stable by definition.

⁶From any infinite sequence starting from $P'[x/y]$ or $P'[* /y]$ we can build an infinite sequence starting from P' .

- If A is a functional type:
 1. Let M be a stable term of type $A \equiv B \rightarrow C$ and let x be a variable of type B . By the second induction hypothesis (with $n = 0$) $x : B$ is stable and so by definition Mx is also stable. By lemma 6.4 Mx is strongly normalizing. Suppose now that M is not strongly normalizing. Then there is an infinite reduction sequence $M \Longrightarrow M_1 \Longrightarrow \dots$ and thus an infinite reduction sequence $Mx \Longrightarrow M_1x \Longrightarrow \dots$ which leads to a contradiction. Therefore M is strongly normalizing.
 2. Let $x\vec{Q}$ be of type $B \rightarrow C$ with all the Q'_i 's strongly normalizing. Let N be any stable term of type B . From the first induction hypothesis N is also strongly normalizing and applying the second induction hypothesis $x\vec{Q}N$ is stable of type C , so by proposition 6.1 $x\vec{Q}$ is stable.

□

Lemma 6.6 *If M is a stable term of type A and $M \Longrightarrow N$, then N is stable.*

Proof. We first recall that N is also of type A by proposition 3.7. We show the property by induction on A .

- A is not a functional type: M is strongly normalizing and then also N is strongly normalizing since for every reduction sequence $N \Longrightarrow^* N'$ there is a longer reduction sequence $M \Longrightarrow N \Longrightarrow^* N'$ that terminates. On the other hand, when $N \Longrightarrow^* \langle M_1, M_2 \rangle$, then M_1 and M_2 are stable, because $M \Longrightarrow^* N \Longrightarrow^* \langle M_1, M_2 \rangle$ and M is stable. Similarly, if $N \Longrightarrow^* in_B^i(M')$ we have $M \Longrightarrow^* N \Longrightarrow in_B^i(M')$ and then M' is stable. Therefore N is stable.
- A is a functional type: by definition of stability on arrow types, it suffices to show that NM_1 is stable for any stable term M_1 . Now, given a stable M_1 , MM_1 is stable because M is, and $MM_1 \Longrightarrow NM_1$, so by induction hypothesis NM_1 is stable.

□

6.3 Products, sums and basic recursion

Lemma 6.7

- $\pi_1(M)$, $\pi_2(M)$ and $in_{A_1+A_2}^i(M)$ are stable if M is stable.
- $Case(P, M_1, M_2)$ is stable if P , M_1 and M_2 are stable.
- $\langle M, N \rangle$ is stable if M and N are stable.
- $(rec\ y : B.M)^0$ is stable if M is stable.

Proof.

- We will prove that $\pi_i(M)$ is stable if M is stable, for $i = 1, 2$. Assume that A is the type of $\pi_i(M)$. Take stable terms \vec{Q} such that $\pi_i(M)\vec{Q}$ is not of functional type. There are two cases to consider:
 - M does not reduce to a pair (recall that M has a product type). By lemma 6.4 $\pi_i(M)\vec{Q}$ is strongly normalizing. On the other hand, $\pi_i(M)\vec{Q}$ cannot reduce to a pair, nor to an in^i because either it reduces to $*$, or the π_i cannot be removed. Therefore $\pi_i(M)\vec{Q}$ is stable for arbitrary stable terms \vec{Q} , so by proposition 6.1 $\pi_i(M)$ is also stable.

- $M \Longrightarrow^* \langle M_1, M_2 \rangle$. Suppose $\pi_i(M)\overrightarrow{Q}$ has an infinite reduction sequence: such reduction has to remove π_i in some step in the following way:

$$\pi_i(M)\overrightarrow{Q} \Longrightarrow^* \pi_i(\langle M_1, M_2 \rangle)\overrightarrow{Q'} \Longrightarrow^* M'_i\overrightarrow{Q''} \Longrightarrow \dots$$

where $M_i \Longrightarrow^* M'_i$ and $Q_i \Longrightarrow^* Q'_i \Longrightarrow^* Q''_i$.

Since M_i and Q_i are stable, by lemma 6.6 also M'_i and Q''_i are stable. Therefore $M'_i\overrightarrow{Q''}$ is stable and then strongly normalizing. From the form of the reduction sequence we can conclude that $\pi_i(M)\overrightarrow{Q}$ is also strongly normalizing which leads to a contradiction. Therefore $\pi_i(M)\overrightarrow{Q}$ has no infinite reduction sequences and is strongly normalizing.

Now, if $\pi_i(M)\overrightarrow{Q}$ reduces to a pair, the reduction removes the π_i and we have necessarily

$$\pi_i(M)\overrightarrow{Q} \Longrightarrow^* \pi_i(\langle M_1, M_2 \rangle)\overrightarrow{Q'} \Longrightarrow^* M'_i\overrightarrow{Q''} \Longrightarrow^* \langle L_1, L_2 \rangle$$

Since $M'_i\overrightarrow{Q''}$ is stable, L_1 and L_2 are stable.

If $\pi_i(M)\overrightarrow{Q}$ reduces to $in^i_C(L)$, the reduction removes the π_i and we have necessarily

$$\pi_i(M)\overrightarrow{Q} \Longrightarrow^* \pi_i(\langle M_1, M_2 \rangle)\overrightarrow{Q'} \Longrightarrow^* M'_i\overrightarrow{Q''} \Longrightarrow^* in^i_C(L)$$

Since $M'_i\overrightarrow{Q''}$ is stable, L is also stable.

We can conclude that $\pi_i(M)\overrightarrow{Q}$ is stable and by proposition 6.1 $\pi_i(M)$ is stable.

- We will prove that $in^i_{A_1+A_2}(M)$ is stable if M is stable. Since M is stable, M is strongly normalizing and by lemma 6.4 also $in^i_{A_1+A_2}(M)$ is strongly normalizing. On the other hand $in^i_{A_1+A_2}(M)$ cannot reduce to a pair because it is of sum type. If $in^i_{A_1+A_2}(M) \Longrightarrow^* in^i_{A_1+A_2}(M')$, then $M \Longrightarrow^* M'$ and since M is stable by hypothesis, we have that M' is stable by lemma 6.6. We conclude that $in^i_{A_1+A_2}(M)$ is stable.
- We will prove that $Case(P, M_1, M_2)$ is stable if P, M_1 and M_2 are stable. Assume that A is the type of $Case(P, M_1, M_2)$. Take stable terms \overrightarrow{Q} such that $Case(P, M_1, M_2)\overrightarrow{Q}$ is not of functional type. Since P is of sum type it cannot reduce to a pair, but it could reduce to an $in^i_B(P')$. There are two cases to consider:
 - P cannot reduce to an in^i . Then by lemma 6.4 $Case(P, M_1, M_2)\overrightarrow{Q}$ is also strongly normalizing. On the other hand, $Case(P, M_1, M_2)\overrightarrow{Q}$ cannot reduce to a pair, nor to an in^i because either $Case(P, M_1, M_2)\overrightarrow{Q}$ reduces to $*$, or the $Case$ construction cannot be removed. Therefore $Case(P, M_1, M_2)\overrightarrow{Q}$ is stable and by proposition 6.1 $Case(P, M_1, M_2)$ is stable.
 - P can reduce to some $in^i_B(P')$. Consider a reduction starting from $Case(P, M_1, M_2)\overrightarrow{Q}$: if in this reduction the $Case(P, M_1, M_2)$ is never eliminated, then the reduction sequence actually proceeds inside the $P, M_1, M_2, \overrightarrow{Q}$, that are all strongly normalizing as they are stable, so this reduction sequence is finite. Otherwise, this reduction sequence removes the $Case$ constructor and looks like:

$$Case(P, M_1, M_2)\overrightarrow{Q} \Longrightarrow^* Case(in^i_B(P'), M'_1, M'_2)\overrightarrow{Q'} \Longrightarrow M'_i P' \overrightarrow{Q''} \Longrightarrow \dots$$

where $M_i \Longrightarrow^* M'_i$ and $Q_i \Longrightarrow^* Q'_i$.

Since P , M_i and the Q_i in \vec{Q} are stable, then P' is stable by definition and by lemma 6.6 also the M'_i and the Q'_i in \vec{Q}' are stable. Therefore $M'_i P' \vec{Q}'$ is stable and then strongly normalizing, so the reduction $M'_i P' \vec{Q}' \Longrightarrow \dots$ is finite and then in turn the whole reduction sequence is finite. We conclude that $Case(P, M_1, M_2) \vec{Q}$ is strongly normalizing. On the other hand, if $Case(P, M_1, M_2) \vec{Q}$ reduces to a pair we have necessarily a reduction sequence removing the *Case* that looks like:

$$Case(P, M_1, M_2) \vec{Q} \Longrightarrow^* Case(in_B^i(P'), M'_1, M'_2) \vec{Q}' \Longrightarrow M'_i P' \vec{Q}' \Longrightarrow^* \langle L_1, L_2 \rangle$$

As seen above, $M'_i P' \vec{Q}'$ is stable, so L_1 and L_2 are stable too.

If $Case(P, M_1, M_2) \vec{Q}$ reduces to $in_C^i(L)$, we have necessarily a reduction sequence removing the *Case* that looks like:

$$Case(P, M_1, M_2) \vec{Q} \Longrightarrow^* Case(in_B^i(P'), M'_1, M'_2) \vec{Q}' \Longrightarrow M'_i P' \vec{Q}' \Longrightarrow^* in_C^i(L)$$

Again, $M'_i P' \vec{Q}'$ is stable, so L is also stable.

We conclude that $Case(P, M_1, M_2) \vec{Q}$ is stable and by proposition 6.1 $Case(P, M_1, M_2)$ is stable.

- We will prove that $\langle M, N \rangle$ is stable if M and N are stable. By hypothesis M and N are strongly normalizing and by lemma 6.4 $\langle M, N \rangle$ is strongly normalizing. It is clear that $\langle M, N \rangle$ cannot reduce to an in^i . If $\langle M, N \rangle \Longrightarrow^* \langle M', N' \rangle$, then $M \Longrightarrow^* M'$ and $N \Longrightarrow^* N'$. By hypothesis M and N are stable and then by lemma 6.6 M' and N' are stable. We can conclude that $\langle M, N \rangle$ is stable.
- We will prove that $(rec\ y : B.M)^0$ is stable if M is stable. Take arbitrary stable terms \vec{Q} such that $(rec\ y : B.M)^0 \vec{Q}$ is not of functional type. By lemma 6.4 $(rec\ y : B.M)^0 \vec{Q}$ is strongly normalizing as the stable terms M , \vec{Q} are strongly normalizing.

On the other hand, the term $(rec\ y : B.M)^0 \vec{Q}$ cannot reduce to a pair, nor to an in^i because either it reduces to $*$, or the *rec* with exponent 0 is not removed. Therefore $(rec\ y : B.M)^0 \vec{Q}$ is stable and by proposition 6.1 $(rec\ y : B.M)^0$ is stable.

□

6.4 Abstraction and recursion

Lemma 6.8 *If $M[N/x]$ is stable, then $(\lambda x : A.M)N$ is stable provided that N is stable if x is not free in M .*

Proof. Let \vec{Q} be stable terms such that $M[N/x] \vec{Q}$ is not of functional type. By proposition 6.1 $M[N/x] \vec{Q}$ is stable and then strongly normalizing. We want to show that $(\lambda x : A.M)N$ is stable, but this follows from proposition 6.1 once we show that $(\lambda x : A.M)N \vec{Q}$ is stable.

Notice first of all that N is strongly normalizing in any case: if x is not free in M we know by hypothesis that N is stable, hence also strongly normalizing, while if x is free in M , then N is strongly normalizing because it is a subterm of the stable (hence strongly normalizing) term $M[N/x]$. Secondly, M is also strongly normalizing, because from a non terminating reduction sequence starting at M we can build a non terminating reduction sequence starting at $M[N/x]$, which is impossible, because this last term is stable, hence strongly normalizing.

We can now show that $(\lambda x : A.M)N\vec{Q}$ is strongly normalizing.

Every reduction sequence starting at $(\lambda x : A.M)N\vec{Q}$ yields to $*$, proceeds inside M , N , \vec{Q} or looks like:

$$(\lambda x : A.M)N\vec{Q} \Longrightarrow^* (\lambda x : A.M')N'\vec{Q}' \Longrightarrow M'[N'/x]\vec{Q}' \Longrightarrow \dots$$

where $M \Longrightarrow M'$, $N \Longrightarrow N'$ and $Q_i \Longrightarrow Q'_i$.

In the first case the reduction stops at $*$, and in the second case the reduction is finite because the terms M , N , \vec{Q} are all strongly normalizing (\vec{Q} are stable, while M and N are strongly normalizing as seen above).

In the last case, by lemma 4.5 and corollary 4.8 $M[N/x] \Longrightarrow^* M'[N'/x]$ and since $M[N/x]$ is stable by hypothesis, applying lemma 6.6 we obtain that $M'[N'/x]$ is also stable. Once again by lemma 6.6 every Q'_i is stable and by proposition 6.1 $M'[N'/x]\vec{Q}'$ is stable and hence strongly normalizing. Then $(\lambda x : A.M)N\vec{Q}$ is also strongly normalizing.

Hence we have proved that $(\lambda x : A.M)N\vec{Q}$ is strongly normalizing.

On the other hand, if $(\lambda x : A.M)N\vec{Q}$ reduces to a pair $\langle L_1, L_2 \rangle$, we have necessarily to remove the redex $(\lambda x : A.M)N$ and we obtain a reduction sequence similar to the previous one with more steps from $M'[N'/x]\vec{Q}'$ to a pair $\langle L_1, L_2 \rangle$. Since $M'[N'/x]\vec{Q}'$ is stable, L_1 and L_2 are stable. Similarly, if $(\lambda x : A.M)N\vec{Q}$ reduces to $in_B^i(L)$, we obtain a reduction sequence with more steps from $M'[N'/x]\vec{Q}'$ to a $in_B^i(L)$. Since $M'[N'/x]\vec{Q}'$ is stable, L is stable.

We can conclude that $(\lambda x : A.M)N\vec{Q}$ is stable and thus we have $(\lambda x : A.M)N$ stable by proposition 6.1. \square

Lemma 6.9 *If $M[(rec\ y.M)^{i-1}/y]$ is stable, then $(rec\ y.M)^i$ is stable.*

Proof. Take stable terms \vec{Q} such that $(rec\ y : B.M)^i\vec{Q}$ is not of functional type. We first show that $(rec\ y : B.M)^i\vec{Q}$ is strongly normalizing. For this, notice that any reduction starting from $(rec\ y : B.M)^i\vec{Q}$ that does not eliminate the rec construct is finite, as it must proceed in the M , \vec{Q} , which are all stable and hence also strongly normalizing. We are left with those reductions that do eliminate the rec construct. Any such reduction sequence looks like:

$$(rec\ y : B.M)^i\vec{Q} \Longrightarrow^* (rec\ y : B.M')^i\vec{Q}' \Longrightarrow M'[(rec\ y : B.M')^{i-1}/y]\vec{Q}'$$

where $Q_i \Longrightarrow^* Q'_i$, $M \Longrightarrow^* M'$ and hence also $M[(rec\ y : B.M)^{i-1}/y] \Longrightarrow^* M'[(rec\ y : B.M')^{i-1}/y]$. Since $M[(rec\ y : B.M)^{i-1}/y]$ and the Q_i are stable, by lemma 6.4 $M'[(rec\ y : B.M')^{i-1}/y]$ and the Q'_i are also stable, so $M'[(rec\ y : B.M')^{i-1}/y]\vec{Q}'$ is stable and thus strongly normalizing. This means that the whole reduction sequence is finite, and concludes this first part of the proof.

Now, if $(rec\ y : B.M)^i\vec{Q}$ reduces to a pair, the rec has been removed and the reduction sequence looks like:

$$(rec\ y : B.M)^i\vec{Q} \Longrightarrow^* (rec\ y : B.M')^i\vec{Q}' \Longrightarrow M'[(rec\ y : B.M')^{i-1}/y]\vec{Q}' \Longrightarrow^* \langle L_1, L_2 \rangle$$

As $M'[(rec\ y : B.M')^{i-1}/y]\vec{Q}'$ is stable, L_1 and L_2 are also stable.

Similarly, if $(rec\ y : B.M)^i\vec{Q}$ reduces to $in_B^i(L)$, the rec has been removed and the reduction sequence looks like:

$$(rec\ y : B.M)^i\vec{Q} \Longrightarrow^* (rec\ y : B.M')^i\vec{Q}' \Longrightarrow M'[(rec\ y : B.M')^{i-1}/y]\vec{Q}' \Longrightarrow^* in_B^i(L)$$

As $M'[(rec\ y : B.M')^{i-1}/y]\vec{Q}'$ is stable, L is also stable.

We can conclude that $(rec\ y : B.M)^i \vec{Q}$ is stable and thus by proposition 6.1 $(rec\ y : B.M)^i$ is stable. \square

Lemma 6.10 *Let M be a term such that all the free variables are among $\{x_i\}_{i=1\dots n}$. If \vec{N} are stable terms, then $M[N_1 \dots N_n/x_1 \dots x_n]$ is stable.*

Proof. We show the property by induction on the structure of M .

- $M \equiv *$. Then $*[N/x_i] = *$ and $*$ is strongly normalizing and hence stable.
- $M \equiv x_i$. Then $x_i[N/x_i] = N$ and N is stable by hypothesis.
- $M \equiv Case(P, Q, R)$. We have $Case(P, Q, R)[\vec{N}/\vec{x}] = Case(P[\vec{N}/\vec{x}], Q[\vec{N}/\vec{x}], R[\vec{N}/\vec{x}])$. By induction hypothesis the terms $P[\vec{N}/\vec{x}]$, $Q[\vec{N}/\vec{x}]$ and $R[\vec{N}/\vec{x}]$ are stable, so we can apply lemma 6.7 and we get that $Case(P[\vec{N}/\vec{x}], Q[\vec{N}/\vec{x}], R[\vec{N}/\vec{x}])$ is stable.
- $M \equiv \pi_i(P)$, for $i = 1, 2$. We have $\pi_i(P)[\vec{N}/\vec{x}] = \pi_i(P[\vec{N}/\vec{x}])$. By induction hypothesis $P[\vec{N}/\vec{x}]$ is stable and by lemma 6.7 $\pi_i(P[\vec{N}/\vec{x}])$ is stable.
- $M \equiv in_B^i(P)$, for $i = 1, 2$. We have $in_B^i(P)[\vec{N}/\vec{x}] = in_B^i(P[\vec{N}/\vec{x}])$. By induction hypothesis $P[\vec{N}/\vec{x}]$ is stable and by lemma 6.7 $in_B^i(P[\vec{N}/\vec{x}])$ is stable.
- $M \equiv \langle M_1, M_2 \rangle$. We have $\langle M_1, M_2 \rangle[\vec{N}/\vec{x}] = \langle M_1[\vec{N}/\vec{x}], M_2[\vec{N}/\vec{x}] \rangle$. By induction hypothesis $M_1[\vec{N}/\vec{x}]$ and $M_2[\vec{N}/\vec{x}]$ are stable and by lemma 6.7 $\langle M_1[\vec{N}/\vec{x}], M_2[\vec{N}/\vec{x}] \rangle$ is stable.
- $M \equiv \lambda y : B.P$. Then $(\lambda y : B.P)[\vec{N}/\vec{x}] = \lambda y : B.P[\vec{N}/\vec{x}]$. Consider any stable term R . By inductive hypothesis $P[\vec{N}/\vec{x}][R/y]$ is stable and by lemma 6.8 $(\lambda y : B.P[\vec{N}/\vec{x}])R$ is stable. By definition of stability $(\lambda y : B.P[\vec{N}/\vec{x}])$ is stable.
- $M \equiv (M_1 M_2)$. We have $(M_1 M_2)[\vec{N}/\vec{x}] = (M_1[\vec{N}/\vec{x}] M_2[\vec{N}/\vec{x}])$. By induction hypothesis $M_1[\vec{N}/\vec{x}]$ and $M_2[\vec{N}/\vec{x}]$ are stable and then by definition of stability for arrow types we conclude that $(M_1[\vec{N}/\vec{x}] M_2[\vec{N}/\vec{x}])$ is stable.
- $M \equiv (rec\ y : B.P)^i$. Then $(rec\ y : B.P)^i[\vec{N}/\vec{x}] = (rec\ y : B.P[\vec{N}/\vec{x}])^i$. By induction hypothesis $P[\vec{N}/\vec{x}][R/y]$ is stable for a stable term R . In particular y is stable and then $P[\vec{N}/\vec{x}][y/y] = P[\vec{N}/\vec{x}]$ is stable.

We will now prove that $(rec\ y : B.P[\vec{N}/\vec{x}])^i$ is stable by induction on i .

- $i = 0$. Since $P[\vec{N}/\vec{x}]$ is stable, this comes from lemma 6.7.
- $i > 0$. Due to lemma 6.9, it suffices to show that $P[\vec{N}/\vec{x}][(rec\ y : B.P[\vec{N}/\vec{x}])^{i-1}/y]$ is stable. But we know by the inductive hypothesis on i that $(rec\ y : B.P[\vec{N}/\vec{x}])^{i-1}$ is stable, and we can then conclude by the inductive hypothesis on the structure.

\square

Corollary 6.11 *Every term is stable, and hence strongly normalizing.*

7 Strong Normalization via reducibility

In this section we will prove the strong normalization property for our calculus $\lambda_{\pi^* \mu \sigma}$, with labeled recursion, but no expansions, using the reducibility method as in [GLT90], with an additional astute twist to take care of the sum type and labeled recursion.

7.1 Reducibility

We define the set RED_A of *reducible* terms of type A by induction on the type A as follows:

- For M of atomic type A , $M \in RED_A$ iff M is strongly normalizable
- For M of product type, $M \in RED_{A_1 \times A_2}$ iff $\pi_i(M) \in RED_{A_i}$
- For M of a sum type, $M \in RED_{A_1 + A_2}$ iff, for fresh variables $w_i : A_i$, we have $Case(M, \lambda x : A_1. \langle x, w_2 \rangle, \lambda y : A_2. \langle w_1, y \rangle) \in RED_{A_1 \times A_2}$ (in the case A_i is \mathbf{T} , we take $*$ instead of w_i)
- For M of a functional type, $M \in RED_{A_1 \rightarrow A_2}$ iff for all $N \in RED_{A_1}$, $(MN) \in RED_{A_2}$

Some comment on the sum type are needed here: first of all notice that the notion of reducibility is well defined: reducibility for a sum type is given in term of reducibility for a product type, which has been defined before. Secondly, notice that for all other types, reducibility is either given directly as in the case of the base types, or given in terms of reducibility for types that are strictly smaller. This is not possible for the sum type, because we have no destructor associated to it, but only a case expression, so reducibility for $A + B$ really depends on reducibility of A and B *together*, and we express this fact by reducing it to reducibility of the product $A \times B$.

7.2 Properties of reducibility

Following [GLT90], we define a notion of *neutrality*: a term is *neutral* if does not interact with the surrounding context giving raise to redexes. In our case, the neutral terms are:

$$* \quad x \quad \pi_i(M) \quad Case(P, M, N) \quad (MN) \quad (rec \ y.M)^i$$

We will prove that RED_A enjoys the following properties, for all types A :

- (CR1) If $M \in RED_A$, then M is strongly normalizable.
- (CR2) If $M \in RED_A$, and M reduces to M' , then $M' \in RED_A$.
- (CR3) If M is neutral and whenever we perform on it one step of reduction we obtain a term $M' \in RED_A$, then $M \in RED_A$.
As a special case of the last clause:
- (CR4) If M is neutral and no reduction is applicable to it, then $M \in RED_A$.

In particular, $*$ and the variables are reducible (also the variables of type \mathbf{T} , as they can only reduce to $*$, which is reducible).

Proposition 7.1 (Properties of reducibility) *For every type A , the set RED_A satisfies (CR1), (CR2) and (CR3).*

Proof. We will proceed by induction on the type A .

7.2.1 Atomic types

- (CR1) A reducible term of atomic type is strongly normalizable by definition
- (CR2) If M is strongly normalizable, then so is every reduct of M (as reduction preserves the type)
- (CR3) Suppose all one step reducts of M are reducible, i.e. strongly normalizable. Any reduction path leaving M must pass through one of its one-step reducts, which are in a finite number, so that the longest reduction sequence starting from M has length the maximum among the $1 + \nu(M')$, as M' varies over the (one-step) reducts of M . Since these lengths are all finite, M is strongly normalizing.

7.2.2 Product types

- (CR1)** Suppose $M \in RED_{A_1 \times A_2}$. Then by definition we know that $\pi_i(M)$ are reducible and so strongly normalizing by induction hypothesis. This implies that M is strongly normalizing also, because any reduction sequence starting from M can be turned into a reduction sequence starting from $\pi_i(M)$.
- (CR2)** We know that $\pi_i(M) \in RED_{A_i}$, by definition. Now consider the possible one step reducts of M :
- M reduces to M' . Then also $\pi_i(M)$ reduces to $\pi_i(M')$ via the same reduction, and M' is then reducible by definition
- (CR3)** Let now M be neutral (not necessarily reducible) such that all its one step reducts are reducible. We must show that $\pi_i(M)$ is reducible of type A_i . Since M cannot be a pair (as it is neutral), any one step reduction of $\pi_i(M)$ must be to a term $\pi_i(M')$, with M' one step from M . By **(CR2)**, M' is reducible, and then by definition also $\pi_i(M')$ is reducible. Now, $\pi_i(M)$ is neutral and all its one step reducts are reducible, hence by induction hypothesis **(CR3)** for A_i , $\pi_i(M)$ is reducible, hence M is, by definition.

7.2.3 Arrow types

- (CR1)** Suppose $M \in RED_{A_1 \rightarrow A_2}$. Then by definition we know that MN is reducible for all reducible N . In particular, Mx is reducible for a fresh variable x , which is reducible by induction hypothesis **(CR3)** for A_1 , hence Mx is strongly normalizable. This implies that also M is strongly normalizing, as all reduction sequences starting from M can be performed also on Mx .
- (CR2)** Let $M \in RED_{A_1 \rightarrow A_2}$ reduce to M' . For all $N \in RED_{A_1}$ we have $(M'N) \in RED_{A_2}$, since it is a reduct of (MN) , which is reducible because M and N are. Hence M' is reducible by definition.
- (CR3)** Let now M be neutral (not necessarily reducible) such that all one step reductions lead to reducible terms. We show that MN is reducible for all reducible N by induction on $\nu(N)$, using **(CR3)** for A_2 . Consider a one step reduction of MN : since M is neutral, this reduction must be either inside M or inside N and leads to:
- $M'N$, with M' one step from M , so M' is reducible and hence $M'N$ is
 - MN' , with N' one step from N ; N' is reducible by **(CR2)** for A_1 , and $\nu(N') < \nu(N)$, so by induction hypothesis MN' is reducible

Hence all reductions leaving MN lead to a reducible term and hence MN is reducible for all reducible N , so that M is reducible by definition.

7.2.4 Sum types

- (CR1)** Suppose $M \in RED_{A_1 + A_2}$. Then by definition $Case(M, \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle) \in RED_{A_1 \times A_2}$ is reducible, hence strongly normalizable, hence M is strongly normalizable too.
- (CR2)** Suppose $M \in RED_{A_1 + A_2}$ reduces to M' . Then $Case(M, \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle) \in RED_{A_1 \times A_2}$ reduces to $Case(M', \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle)$, so that, by **(CR2)** for $A \times B$ which has been proved before, $Case(M, \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle) \in RED_{A_1 \times A_2}$. Hence M' is reducible too.
- (CR3)** Let now M be neutral, and suppose all its one step reducts are reducible. We will show that $Case(M, \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle)$ (which is neutral) is reducible using **(CR3)** for $A_1 \times A_2$, which has already been proved to hold. Consider the possible one step reducts:
- $Case(M', \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle)$ with M' one step from M : then M' is reducible, hence $Case(M', \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle)$ is reducible by definition

- there is no other one step reduct as M is neutral and the terms $\lambda x.\langle x, w_2 \rangle$ and $\lambda y.\langle w_1, y \rangle$ are normal

□

7.3 Reducibility theorem

We are left to show a few more lemmas:

Lemma 7.2 (Pairing) *Let $M_1 : A_1, M_2 : A_2$ be reducible terms. Then $\langle M_1, M_2 \rangle \in RED_{A_1 \times A_2}$.*

Proof. We need to show that $\pi_i(\langle M_1, M_2 \rangle) \in RED_{A_i}$.

Since $\pi_i(\langle M_1, M_2 \rangle)$ is neutral, we prove the statement using **(CR3)**: we will show that all one step reductions are reducible. We proceed by induction on the sum $\nu(M_1) + \nu(M_2)$ of the maximum reduction lengths for M_1 and M_2 , (which are finite, as these terms are strongly normalizable by **(CR1)**).

The possible reducts are:

- M_i , which is reducible by hypothesis
- $\pi_i(\langle M'_1, M_2 \rangle)$: now, M'_1 is one step from M_1 , so that $\nu(M'_1) + \nu(M_2) < \nu(M_1) + \nu(M_2)$, and M'_1 is reducible by **(CR2)**, so $\pi_i(\langle M'_1, M_2 \rangle)$ is reducible by induction hypothesis
- $\pi_i(\langle M_1, M'_2 \rangle)$: this is shown reducible as the term in the previous case

□

Lemma 7.3 (Abstraction) *Let $M : A_2$ be a term where the variable $x : A_1$ may occur free. If for every $N \in RED_{A_1}$ we have $M[N/x] \in RED_{A_2}$, then $\lambda x : A_1.M \in RED_{A_1 \rightarrow A_2}$.*

Proof. We want to show that $(\lambda x.M)P$ is reducible for all reducible P . Since this term is neutral, we can prove our Lemma using **(CR3)**. We are then left to show that all one step reducts of $(\lambda x.M)P$ are reducible if for all $N \in RED_{A_1}$ we have $M[N/x] \in RED_{A_2}$. Since $M = M[x/x]$ is reducible by hypothesis (as any variable is reducible), it is strongly normalizable by **(CR1)**, and we can proceed to prove this last statement by induction on $\nu(M) + \nu(P)$. The term $(\lambda x.M)P$ converts to:

- $M[P/x]$ which is reducible by hypothesis
- $(\lambda x.M')P$ with M' a reduct of M ; now, by **(CR2)**, M' is still reducible and furthermore $\nu(M') + \nu(P) < \nu(M) + \nu(P)$ and $M[P/x]$ reduces to $M'[P/x]$, and this last term is also reducible, because it is a multi-step reduct of $M[P/x]$ by Lemma 4.5. So the induction hypothesis tells us that $(\lambda x.M')P$ is reducible.
- $(\lambda x.M)P'$ with P' a reduct of P ; now, by **(CR2)**, P' is still reducible and furthermore $\nu(M) + \nu(P') < \nu(M) + \nu(P)$ and $M[P/x]$ reduces to $M[P'/x]$, by Corollary 4.8, so this last term is also reducible. The induction hypothesis tells us that $(\lambda x.M)P'$ is reducible.

□

Lemma 7.4 (Injections) *For all terms, $M \in RED_{A_1}$ iff $in^i_{A_1+A_2}(M) \in RED_{A_1+A_2}$.*

Proof. (\Rightarrow)

We must show that $Case(in^i_{A_1+A_2}(M), \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle)$ is reducible of type $A_1 \times A_2$, i.e. that $\pi_i(Case(in^i_{A_1+A_2}(M), \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle)) \in RED_{A_i}$. Since $\pi_i(Case(in^i_{A_1+A_2}(M), \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle))$ is neutral, we will proceed using **(CR3)**, by induction on $\nu(M)$.

Consider then all its one step reducts:

- $((\lambda x.\langle x, w_2 \rangle)M)$, which is reducible because M is reducible and $\lambda x.\langle x, w_2 \rangle$ is reducible (by Lemma 7.2 applied to the variables x and w_2 we know that $\langle x, w_2 \rangle$ is reducible, and we get reducibility of $\lambda x.\langle x, w_2 \rangle$ by Lemma 7.3; similarly if we have $*$ instead of w_2)

- $\pi_i(\text{Case}(\text{in}_{A_1+A_2}^i(M'), \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle))$ with M' one step from M , hence reducible and $\nu(M') < \nu(M)$, so that it is reducible by induction hypothesis

(\Leftarrow)

Suppose now, $\text{in}_{A_1+A_2}^i(M) \in \text{RED}_{A_1+A_2}$. This means, by definition of reducibility over sum types,

$$\text{Case}(\text{in}_{A_1+A_2}^i(M), \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle) \in \text{RED}_{A_1 \times A_2},$$

which implies, by definition of reducibility over product types,

$$\pi_1(\text{Case}(\text{in}_{A_1+A_2}^i(M), \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle)) \in \text{RED}_{A_1}$$

This term reduces to $\pi_1((\lambda x.\langle x, w_2 \rangle)M)$, which in turn reduces to $\pi_1(\langle M, w \rangle)$ and then to M , which is then reducible by repeated use of **(CR2)**. \square

Lemma 7.5 (Sum) *Let $P : A + B$, $M : A \rightarrow C$ and $N : B \rightarrow C$ be reducible terms. Then $\text{Case}(P, M, N) \in \text{RED}_C$.*

Proof. We will work by cases on C .

C is an atomic type We can use **(CR3)** for C , as $\text{Case}(P, M, N)$ is neutral. We will show by induction on $\nu(P) + \nu(M) + \nu(N)$ that all one step reducts of $\text{Case}(P, M, N)$ are reducible. Consider the possible one step reducts:

- $\text{Case}(P', M, N)$, or $\text{Case}(P, M', N)$, or $\text{Case}(P, M, N')$: they are reducible by induction hypothesis as all primed terms are reducible by **(CR2)** on $A + B$, $A \rightarrow C$, $B \rightarrow C$, and the measure decreases strictly.
- (RM) if $P \equiv \text{in}_{A_1+A_2}^i(R)$: then R is also reducible by Lemma 7.4, and this term is reducible as M is

$C \equiv C_1 \times C_2$ We must show $\pi_i(\text{Case}(P, M, N)) \in \text{RED}_{C_i}$. Since $\pi_i(\text{Case}(P, M, N))$ is neutral, we can use **(CR3)** for C_i . Since P, M, N , are all reducible, they are all strongly normalizable and we can proceed by induction on the measure $\nu(P) + \nu(M) + \nu(N)$. Consider the possible one step reducts:

- $\pi_i(\text{Case}(P', M, N))$ or $\pi_i(\text{Case}(P, M', N))$ or $\pi_i(\text{Case}(P, M, N'))$: they are reducible by induction hypothesis as all primed terms are reducible by **(CR2)** on $A + B$, $A \rightarrow C$, $B \rightarrow C$, and the measure decreases strictly.
- $\pi_i((MR))$ if $P \equiv \text{in}_{A_1+A_2}^i(R)$: then R is also reducible by Lemma 7.4, so MR is reducible and $\pi_i((MR))$ too

$C \equiv C_1 \rightarrow C_2$ We must show $\text{Case}(P, M, N)Q \in \text{RED}_{C_2}$ for all $Q \in \text{RED}_{C_1}$. Since $\text{Case}(P, M, N)Q$ is neutral, we can use **(CR3)** for C_2 . Since P, M, N, Q are all reducible, they are all strongly normalizable and we can proceed by induction on the measure $\nu(P) + \nu(M) + \nu(N) + \nu(Q)$. Consider the possible one step reducts:

- $\text{Case}(P', M, N)Q$, or $\text{Case}(P, M', N)Q$, or $\text{Case}(P, M, N')Q$, or $\text{Case}(P, M, N)Q'$: they are reducible by induction hypothesis as all primed terms are reducible by **(CR2)** on $A+B$, $A \rightarrow C$, $B \rightarrow C$ and C_1 , and the measure decreases strictly.
- $(RM)Q$ if $P \equiv \text{in}_{A_1+A_2}^i(R)$: then R is also reducible by Lemma 7.4, and this term is reducible as M and Q are

$C \equiv C_1 + C_2$ We must show $\text{Case}(\text{Case}(P, M, N), \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle) \in \text{RED}_{C_1 \times C_2}$. We can use **(CR3)** for $C_1 \times C_2$ because $\text{Case}(\text{Case}(P, M, N), \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle)$ is neutral. Since P, M, N , are all reducible, they are all strongly normalizable and we can proceed by induction on the measure $\nu(P) + \nu(M) + \nu(N)$. Consider the possible one step reducts:

- $Case(Case(P', M, N), \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle)$ or $Case(Case(P, M', N), \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle)$ or $Case(Case(P, M, N'), \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle)$: they are reducible by induction hypothesis as all primed terms are reducible by **(CR2)** on $A + B$, $A \rightarrow C$, $B \rightarrow C$, and the measure decreases strictly.
- $Case((RM), \lambda x.\langle x, w_2 \rangle, \lambda y.\langle w_1, y \rangle)$ if $P \equiv in_{A_1+A_2}^i(R)$: then R is also reducible by Lemma 7.4, and this term is reducible by definition as M , hence also (RM) , is

□

We will now prove that every reducible instance of a (not necessarily reducible) term M is reducible. As a consequence, all terms will be reducible.

Theorem 7.6 (Reducibility) *Let M be any term (not assumed to be reducible), and suppose all the free variables of M are among x_1, \dots, x_n of types A_1, \dots, A_n . If N_1, \dots, N_n are reducible terms of types A_1, \dots, A_n , then $M[\overline{N}/\overline{x}]$ is reducible.*

Proof. By induction on the structure of M .

1. M is $*$. It is neutral and normal, so it is reducible.
2. M is x_i for some i , then $M[\overline{N}/\overline{x}] = N_i$ is reducible
3. $M \equiv \pi_i(M')$. By induction hypothesis, $M'[\overline{N}/\overline{x}]$ is reducible, hence, by definition, $\pi_i(M'[\overline{N}/\overline{x}]) = \pi_i(M')[\overline{N}/\overline{x}]$ is reducible.
4. $M \equiv \langle M_1, M_2 \rangle$. By induction hypothesis, the terms $M_i[\overline{N}/\overline{x}]$ are reducible, so we conclude that the term $\langle M_1[\overline{N}/\overline{x}], M_2[\overline{N}/\overline{x}] \rangle = \langle M_1, M_2 \rangle[\overline{N}/\overline{x}]$ is reducible.
5. $M \equiv in_{A_1+A_2}^i(M')$. By induction hypothesis, $M'[\overline{N}/\overline{x}]$ is reducible, hence by Lemma 7.4, $in_{A_1+A_2}^i(M'[\overline{N}/\overline{x}]) = in_{A_1+A_2}^i(M')[\overline{N}/\overline{x}]$ is reducible.
6. $M \equiv Case(M_1, M_2, M_3)$. By induction hypothesis, the terms $M_i[\overline{N}/\overline{x}]$ are reducible, hence $Case(M_1[\overline{N}/\overline{x}], M_2[\overline{N}/\overline{x}], M_3[\overline{N}/\overline{x}]) = Case(M_1, M_2, M_3)[\overline{N}/\overline{x}]$ is reducible.
7. $M \equiv (M_1 M_2)$. By induction hypothesis, the terms $M_i[\overline{N}/\overline{x}]$ are reducible, so we conclude that the term $(M_1[\overline{N}/\overline{x}])M_2[\overline{N}/\overline{x}] = (M_1 M_2)[\overline{N}/\overline{x}]$ is reducible.
8. $M \equiv \lambda y.M'$. Then by induction hypothesis $M'[\overline{N}/\overline{x}][N'/y]$ is reducible for all reducible terms N' . By Lemma 7.3, $\lambda y.M'[\overline{N}/\overline{x}] = (\lambda y.M')[\overline{N}/\overline{x}]$ is reducible.
9. $M \equiv (recy.M')^i$. By induction hypothesis, $M'[\overline{N}/\overline{x}]$ is reducible. We will show reducibility for $(recy.M')^i[\overline{N}/\overline{x}]$ by induction on $i + \nu(M')$. Since $(recy.M)^i[\overline{N}/\overline{x}]$ is neutral, we will use **(CR3)** for the type A of $(recy.M)^i$. Consider the one step reducts of $(recy.M)^i[\overline{N}/\overline{x}]$
 - $(recy.M'')^i[\overline{N}/\overline{x}]$ with M'' one step from M' . Then $M''[\overline{N}/\overline{x}]$ is reducible for all reducible \overline{N} , because it is a multi-step reduct of the reducible term $M'[\overline{N}/\overline{x}]$ (Lemma 4.5). Furthermore, $i + \nu(M'') < i + \nu(M')$, so by induction hypothesis $(recy.M'')^i[\overline{N}/\overline{x}] = (recy.M'')^i[\overline{N}/\overline{x}]$ is reducible.
 - $M'[\overline{N}/\overline{x}][(recy.M'[\overline{N}/\overline{x}])^{i-1}/y]$. Then $(recy.M')^{i-1}[\overline{N}/\overline{x}] = (recy.M'[\overline{N}/\overline{x}])^{i-1}$ is reducible by induction hypothesis, and this tells us that $[\overline{N}/\overline{x}][(recy.M'[\overline{N}/\overline{x}])^{i-1}/y]$ is a substitution of reducible terms for a set of variables containing the free variables of M' , which gives us reducibility of the term $M'[\overline{N}/\overline{x}][(recy.M'[\overline{N}/\overline{x}])^{i-1}/y]$.

□

Corollary 7.7 (Strong Normalization) *All terms are reducible, hence strongly normalizable.*

8 Confluence of the Full Calculus

In this section we deduce the confluence property for the calculus with bounded recursion as well as for the version with unbounded recursion.


We can immediately deduce the confluence property for the bounded system from the weak confluence and strong normalization properties using Newman's Lemma, however, we can also provide an extremely simple and neat proof that does not need the weak confluence property for the expansionary system.

Theorem 8.1 (Confluence) *The relation \Longrightarrow is Church-Rosser.*

Proof. Since \Longrightarrow is weakly confluent by theorem 4.11 and strongly normalizing by theorem 5.14 we can conclude that it is Church Rosser by the well known Newman's lemma.

The other proof of confluence proceeds as follows.

Let M be a term s.t. $P_1 \ast \leftarrow M \Longrightarrow^* P_2$. Since \Longrightarrow is strongly normalizing, we can reduce the terms P_i to their normal forms \overline{P}_i . Then we have $\overline{P}_1 \ast \leftarrow M \Longrightarrow^* \overline{P}_2$, and by theorem 5.13 $\overline{P}_1^\circ \ast \leftarrow M^\circ \Longrightarrow^* \overline{P}_2^\circ$ without expansions in the reduction sequences. As the system without expansions is confluent (we showed that it is strongly normalizing, and weak confluence without expansions can be shown as easily as for the simply typed lambda calculus), we can close the internal diagram with $\overline{P}_1^\circ \Longrightarrow^* R \ast \leftarrow \overline{P}_2^\circ$. Now, $\overline{P}_i^\circ =_{\text{lemma 5.6}} \overline{P}_i$ and therefore we can complete the proof using the reductions $P_1 \Longrightarrow^* \overline{P}_1 \Longrightarrow^* R \ast \leftarrow \overline{P}_2 \ast \leftarrow P_2$ (notice that $\overline{P}_1 = R = \overline{P}_2$). The following figure shows the reduction diagram:



□

In order to show confluence of the full calculus we relate in the first place the bounded reduction \Longrightarrow and the unbounded one \Longrightarrow^∞ , and then we use the confluence of \Longrightarrow to show the confluence of \Longrightarrow^∞ . This very same technique, that originates from early work of Lévy [Lév76], was used in [PV87]. The connection between the reductions \Longrightarrow and \Longrightarrow^∞ comes from the following:

Remark 8.2 *If $M \Longrightarrow^* N$, then $|M| \Longrightarrow^\infty |N|$, where $|M|$ is obtained from M by removing all the indices from the rec terms.*

Lemma 8.3 *For any reduction sequence $M_0 \Longrightarrow^\infty M_1 \Longrightarrow^\infty \dots \Longrightarrow^\infty M_n$, there exists an indexed computation $N_0 \Longrightarrow N_1 \Longrightarrow \dots \Longrightarrow N_n$ such that $|N_i| = M_i$, for $i = 0 \dots n$.*

Proof. Index all the rec constructors in M_0 by a number $n + k$, with $k \geq 0$. □

Confluence of the full calculus results now from the confluence of the bounded calculus.

Theorem 8.4 \Longrightarrow^∞ is Church Rosser.

Proof. Let $M \equiv P_0 \xRightarrow{\infty} P_1 \xRightarrow{\infty} \dots \xRightarrow{\infty} P_n$ and $M \equiv Q_0 \xRightarrow{\infty} Q_1 \xRightarrow{\infty} \dots \xRightarrow{\infty} Q_m$.

By lemma 8.3 there are indexed computations

$$P'_0 \xRightarrow{\infty} P'_1 \xRightarrow{\infty} \dots \xRightarrow{\infty} P'_n \text{ and } Q'_0 \xRightarrow{\infty} Q'_1 \xRightarrow{\infty} \dots \xRightarrow{\infty} Q'_m$$

where $|P'_i| = P_i$, for $i = 0 \dots n$ and $|Q'_i| = Q_i$, for $i = 0 \dots m$.

We can assume that $P'_0 \equiv Q'_0$ by indexing P^0 and Q^0 with $n + m$. As $\xRightarrow{\infty}$ is Church Rosser by theorem 8.1, there exists a term R such that $P'_n \xRightarrow{\infty} R$ and $Q'_m \xRightarrow{\infty} R$. By 8.2 $P_n = |P'_n| \xRightarrow{\infty} R$ and $Q_m = |Q'_m| \xRightarrow{\infty} R$. \square

9 Adding weak extensionality for the sum type

In this section we show how to apply our techniques in order to accomodate in our system the weak extensionality for the sum type, that is described by the following equality, which tells us that any term P of sum type $A_1 + A_2$ is definitely an injection from one of the two types A_i .

$$\text{Case}(P, \lambda x.in^1(x), \lambda y.in^2(y)) = P \tag{1}$$

This is the usual equality that is found in proof theory, associated to the logical connective for disjunction (see for example [GLT90, Gir72]). We call this rule “weak” because in category theory there is another stronger kind of extensional equality associated to the sum, that is used to axiomatize the uniqueness of the sum of two arrows in the diagram for the coproduct, namely

$$\text{Case}(P, M \circ \lambda x.in^1(x), M \circ \lambda y.in^2(y)) = MP \tag{2}$$

where $M \circ N$ is the usual abbreviation for the composition $\lambda x.(M(Nx))$.

One can easily see that this strong rule really breaks down into two simpler rules: the weak rule 1 we just introduced and the following commutation rule:

$$\text{Case}(P, M \circ N_1, M \circ N_2) = M\text{Case}(P, N_1, N_2) \tag{3}$$

If one really wants the equality 2, it seems to be a difficult task to provide a confluent system for the extensional theory with arrow, product and coproduct types, as discussed in [Dou90], and to the author’s best knowledge, there are no positive results in that direction.

Notice also that the equation 1 can be easily added to a reduction system with no \mathbf{T} type, where all the extensional equalities are turned into *contractions*, as done for example in [Gal93]. In the presence of the \mathbf{T} type, to use contraction rules one is forced to proceed along the lines of [CDC91], and to generate an infinite set of reduction rules.

It is not obvious to add weak extensionality for the sum to our system, as the naïve idea of adding the equality 1 as a contraction rule breaks confluence, as the following example shows:

$$\begin{array}{c} \text{Case}(w, \lambda x : A \rightarrow B.in^1_{(A \rightarrow B)+C}(x), \lambda y : C.in^2_{(A \rightarrow B)+C}(y)) \longrightarrow w \\ \Downarrow \\ \text{Case}(w, \lambda x : A \rightarrow B.in^1_{(A \rightarrow B)+C}(\lambda z : A.xz), \lambda y : C.in^2_{(A \rightarrow B)+C}(y)) \end{array}$$

This problem comes from the fact that the term $\lambda x : A \rightarrow B.in^1_{(A \rightarrow B)+C}(x)$ is not in normal form w.r.t. the rules η , δ and Top . This also suggests the solution: it suffices to completely expand the terms $IN^1 = \lambda x.in^1(x)$ and $IN^2 = \lambda y.in^2(y)$ w.r.t. the rules η , δ and Top (which we know now are strongly normalizing) before performing the contraction for the weak sum extensional equality.

So we are led to consider the contraction rule:

$$\text{Case}(P, \|\!|IN^1\|\!, \|\!|IN^2\|\!) \xrightarrow{+} P$$

where $\|\!|M\|\!$ denotes the normal form of M w.r.t. η , δ and Top expansions.

It is now straightforward to check that the weak confluence property still holds, and one is left to check that the simulation theorem stays valid.

For that, we have to verify that $Case(P, \|IN^1\|, \|IN^2\|)^\circ \Longrightarrow^+ P^\circ$ without using expansion rules in the reduction sequence, and this is obtained by:

$$\begin{aligned} Case(P, \|IN^1\|, \|IN^2\|)^\circ &= \\ \Delta_{A+B}^k Case(P^\circ, \|IN^1\|^\circ, \|IN^2\|^\circ) &= \\ \Delta_{A+B}^k Case(P^\circ, \|IN^1\|, \|IN^2\|) &\xrightarrow{+} \\ \Delta_{A+B}^k P^\circ &\Longrightarrow^* \\ P^\circ & \end{aligned}$$

Notice that the rules η , δ and Top do not create new redexes, as shown in lemma 3.8, so in particular $\|IN^i\|$ is still in normal form, and the equality $\|IN^i\|^\circ = \|IN^i\|$ can be obtained from lemma 5.6.

10 Conclusion and Future Work

We have provided a confluent rewriting system for an extensional typed λ -calculus with product, sum, terminal object and recursion, which is also strongly normalizing in case the recursion operator is bounded. There are mainly two relevant technical contributions in this paper: the weak confluence proof and the simulation theorem.

On one hand, let us remark once again that the weak confluence property for a context-sensitive reduction system is not as straightforward as for the reduction systems that are congruencies. The proof is no longer just a matter of a boring but trivial case analysis, so we had to explore and analyze here the fine structure of the reduction system, showing clearly how substitution and reduction interact in the presence of context-sensitive rules.

The simulation theorem, on the other hand, turns out to be the real key tool for this expansionary system: it allows to reduce *both* confluence *and* strong normalization properties to those for the underlying calculus without expansions, that can be proved using the standard techniques. In a sense, this is all that you really need to prove.

References

- [Aka93] Y. Akama. On mints' reductions for ccc-calculus. In *Typed Lambda Calculus and Applications*, number 664 in LNCS. Springer Verlag, 1993.
- [Bar84] H. Barendregt. *The Lambda Calculus; Its syntax and Semantics (revised edition)*. North Holland, 1984.
- [CDC91] P.-L. Curien and R. Di Cosmo. A confluent reduction system for the λ -calculus with surjective pairing and terminal object. In Leach, Monien, and Artalejo, editors, *Intern. Conf. on Automata, Languages and Programming (ICALP)*, number 510 in LNCS, pages 291–302. Springer-Verlag, 1991.
- [CF58] H. Curry and R. Feys. *Combinatory Logic*, volume 1. North Holland, 1958.
- [Cub92] D. Cubric. On free ccc. Distributed on the `types` mailing list, 1992.
- [DCK93] R. Di Cosmo and D. Kesner. A confluent reduction for the extensional typed λ -calculus with pairs, sums, recursion and terminal object. In A. Lingas, editor, *Intern. Conf. on Automata, Languages and Programming (ICALP)*, LNCS. Springer-Verlag, 1993.
- [Dou90] D. J. Dougherty. Some reduction properties of a lambda calculus with coproducts and recursive types. Technical report, Wesleyan University, 1990. E-mail: `ddougherty@eagle.wesleyan.edu`.
- [Dou93] D. J. Dougherty. Some lambda calculi with categorical sums and products. In *Proc. of the Fifth International Conference on Rewriting Techniques and Applications (RTA)*, 1993.

- [Gal93] J. Gallier. On the correspondence between proofs and λ -terms. Available by anonymous ftp from ftp.cis.upenn.edu. Author's address: jean@saul.cis.upenn.edu., January 1993.
- [Gir72] J.-Y. Girard. *Interprétation fonctionnelle et élimination des coupures dans l'arithmétique d'ordre supérieure*. Thèse de doctorat d'état, Université de Paris VII, 1972.
- [GLT90] J.-Y. Girard, Y. Lafont, and P. Taylor. *Proofs and Types*. Cambridge University Press, 1990.
- [Hue76] G. Huet. Résolution d'équations dans les langages d'ordre $1, 2, \dots, \omega$. *Thèse d'Etat, Université Paris VII*, 1976.
- [Jay92] C. B. Jay. Long $\beta\eta$ normal forms and confluence (and its revised version). Technical Report ECS-LFCS-91-183, LFCS, University of Edimburgh, 1992.
- [JG92] C. B. Jay and N. Ghani. The virtues of eta-expansion. Technical Report ECS-LFCS-92-243, LFCS, University of Edimburgh, 1992.
- [Klo80] J. W. Klop. Combinatory reduction systems. *Mathematical Center Tracts*, 27, 1980.
- [Kri90] J.-L. Krivine. *Lambda calculus. Types et Modèles*. Masson, 1990.
- [Lév76] J.-J. Lévy. An algebraic interpretation of the $\lambda\beta\kappa$ -calculus and a labelled λ -calculus. *Theoretical Computer Science*, 2:97–114, 1976.
- [LS86] J. Lambek and P. J. Scott. *An introduction to higher order categorical logic*. Cambridge University Press, 1986.
- [Min77] G. Mints. Closed categories and the theory of proofs. *Zapiski Nauchnykh Seminarov Leningradskogo Otdeleniya Matematicheskogo Instituta im. V.A. Steklova AN SSSR*, 68:83–114, 1977.
- [Min79] G. Mints. Teorija categorii i teoria dokazatelstv.I. *Aktualnye problemy logiki i metodologii nauky*, pages 252–278, 1979.
- [Pot81] G. Pottinger. The Church Rosser Theorem for the Typed lambda-calculus with Surjective Pairing. *Notre Dame Journal of Formal Logic*, 22(3):264–268, 1981.
- [Pra71] D. Prawitz. Ideas and results in proof theory. *Proceedings of the 2nd Scandinavian Logic Symposium*, pages 235–30. , 1979.
- [PV8.433749(7)0.433749(9)] A. Poigné and J. Voss. On the implementation of typed lambda programming language constructs. *Journal of Computer and System Science*, 34(2-3):340–376, April/June 1987.
- [Tro86] A. S. Troelstra. Strong normalization for typed terms with surjective pairing. *Notre Dame Journal of Formal Logic*, 27(4), 1986.